


Figure Chromatograms illustrating a novel *ATP1A3* gene mutation in a child with rapid-onset dystonia-parkinsonism



Sequence from the patient and a control. Chromatograms show the mutation found in the patient in the *ATP1A3* gene. (A) Wild type sequence. The arrow in (B) shows the heterozygous mutant sequence at position c.2767 G>A that results in a D923N amino acid substitution in the protein.

the membrane and lies close to the ion-binding residue Q920. Prior mutations of Na,K-ATPase provide insight into the D923N mutation. In α_2 , mutations of the equivalent residue, D925L or D925N, had residual Na,K-ATPase activity; however, affinities for both Na^+ and K^+ were significantly reduced.⁶ In α_1 subunit, mutation of the glutamine equivalent to Q920 in the human α_3 subunit gene abolished activity, consistent with its role in ion binding.⁷ Thus D923 is close enough to the ion-binding cavity to affect enzyme activity when mutated. Based on the behavior of the same substitution in the α_2 subunit, we expect Na,K-ATPase in patients with this mutation to have impaired activity and ion-binding affinities.

The novel mutation detected in our patient seems to be a causative one. This patient's presentation has distinctive features and demonstrates that early childhood onset of RDP may occur.

From the Department of Neurology (I.A.A., B.T.D.), Children's Hospital Boston and Harvard Medical School; Department of Surgery (K.J.S.), Massachusetts General Hospital and Harvard Medical School, Boston, MA; and Department of Genetics and Genomic Sciences and Department of Neurology (S.G., L.J.O.), Mount Sinai School of Medicine, New York, NY.

Dr. Sweadner was supported by the Bachmann-Strauss Dystonia and Parkinson Foundation. Drs. Sweadner and Ozelius were supported by NIH grant NS058949.

Disclosure: The authors report no disclosures.

Received October 17, 2008. Accepted in final form March 13, 2009.

Address correspondence and reprint requests to Dr. Basil T. Darras, Department of Neurology, Fegan 11, Children's Hospital Boston, 300 Longwood Avenue, Boston, MA 02115; basil.darras@childrens.harvard.edu

Copyright © 2009 by AAN Enterprises, Inc.

1. de Carvalho Aguiar P, Sweadner KJ, Penniston JT, et al. Mutations in the Na⁺/K⁺-ATPase alpha3 gene *ATP1A3* are associated with rapid-onset dystonia parkinsonism. *Neuron* 2004;43:169–175.
2. Brashear A, Dobyns WB, de Carvalho Aguiar P, et al. The phenotypic spectrum of rapid-onset dystonia-parkinsonism (RDP) and mutations in the *ATP1A3* gene. *Brain* 2007; 130(Pt 3):828–835.
3. Brashear A, Farlow MR, Butler IJ, Kasarskis EJ, Dobyns WB. Variable phenotype of rapid-onset dystonia-parkinsonism. *Mov Disord* 1996;11:151–156.
4. Pittcock SJ, Joyce C, O'Keane V, et al. Rapid-onset dystonia-parkinsonism: a clinical and genetic analysis of a new kindred. *Neurology* 2000;55:991–995.
5. Morth JP, Pedersen BP, Toustrup-Jensen MS, et al. Crystal structure of the sodium-potassium pump. *Nature* 2007; 450:1043–1049.
6. Yamamoto S, Kuntzweiler TA, Wallick ET, Sperelakis N, Yatani A. Amino acid substitutions in the rat Na⁺, K(+) -ATPase alpha 2-subunit alter the cation regulation of pump current expressed in HeLa cells. *J Physiol* 1996;495(Pt 3): 733–742.
7. Arguello JM, Whitis J, Lingrel JB. Alanine scanning mutagenesis of oxygen-containing amino acids in the transmembrane region of the Na,K-ATPase. *Arch Biochem Biophys* 1999;367:341–347.

Stephen Tisch, MBBS,
PhD
Bruce Brew, MBBS, MD

PARKINSONISM IN HIV-INFECTED PATIENTS ON HIGHLY ACTIVE ANTIRETROVIRAL THERAPY

Highly active antiretroviral therapy (HAART) has significantly prolonged the lives of HIV-infected patients but low level chronic immune activation remains in some. Neurodegenerative diseases are often associated with immune activation which likely plays a role in their pathogenesis.¹ Recently, increased alpha-synuclein deposition in the substantia nigra in aging HIV-infected patients has been identified,²

although there was no clear clinical correlate. These observations suggest that HIV and HAART may facilitate the development of neurodegeneration and accelerate the appearance of diseases such as Parkinson disease (PD) via mechanisms including inflammation, mitochondrial dysfunction, and interference with the ubiquitin proteasome pathway.¹ Here we describe 3 patients with relatively young onset of parkinsonism in the context of HIV and HAART.

Table	Patient clinical data		
	Patient 1	Patient 2	Patient 3
Gender	Male	Male	Male
Age, y	55	44	53
Duration of HIV, y	10	4	13
Duration of HAART, y	3	4	10
Previous HAART	Efavirenz, emtricitabine, tenofovir	Ritonavir, indinavir, lamivudine, abacavir, atazanavir, efavirenz	Zidovudine, lamivudine, nelfinavir, nevirapine, ritonavir, saquinavir, atazanavir, didanosine
Current HAART	Nevirapine, emtricitabine, tenofovir	Raltegravir, abacavir, lamivudine	Efavirenz, emtricitabine, tenofovir
Current HIV viral load	<50 copies/mL	<50 copies/mL	<50 copies/mL
Nadir and current CD4 count	180/380	510/728	88/450
Duration of parkinsonism, y	2	3	3
Initial symptoms	Rest tremor of the left hand	Rest tremor of the right hand	Rest tremor of the right hand
Clinical features	Hypomimia, monotonous speech, left-sided rest tremor, rigidity and bradykinesia, loss of left arm swing	Rest tremor of the right hand, mild symmetric limb rigidity, bradykinesia, bilaterally reduced arm swing, foot dystonia	Hypomimia, right-sided rest tremor, rigidity, bradykinesia, bilaterally reduced arm swing, REM sleep behavior disorder
Hoehn and Yahr score (1-5)	1	2	2
Cognitive impairment	No	No	No
L-dopa response (max dose)	Nil (400 mg/d)	Nil (600 mg/d)	Good (300 mg/d)
Brain MRI	Spectroscopy reduced NAA/creatine in caudate, otherwise normal	Spectroscopy reduced NAA/creatine in caudate, otherwise normal	Normal
CSF	Not sampled	No cells, protein 906 mg/L, neopterin 40 (0-13 nmol/L), β 2 microglobulin 2.3 (<2.2 mg/L), HIV viral load 100 copies/mL	Normal protein, glucose, cell count, β 2 microglobulin, neopterin HIV viral load 230 copies/mL

HAART = highly active antiretroviral therapy.

Case series. The patients were HIV-infected men aged between 44 and 55 years on HAART who were examined prior to the development of parkinsonism. None had been exposed to neuroleptic medications or recent amphetamine use. All developed initial unilateral upper extremity rest tremor, followed by rigidity, bradykinesia, and mild gait involvement with preserved postural stability. Importantly, cognitive, oculomotor, pyramidal, cerebellar, and autonomic features were absent. Only patient 3 responded well to levodopa; however, the dose in patients 1 and 2 could not be raised due to side effects. In all patients, parkinsonism developed and progressed in the context of effective HAART (plasma HIV viral load <50 copies/mL). Brain MRI was normal apart from reduced *N*-acetylaspartate spectroscopy peaks (relative to creatine) in the caudate in 2 patients. In 2 patients, CSF showed low levels of HIV RNA and 1 had evidence of inflammation with raised neopterin and β 2 microglobulin. The patient details are described in the table.

Discussion. HAART has transformed HIV to a chronic disease with near-normal life expectancy. There is accumulating evidence that HIV and HAART accelerate the aging process, thereby possibly promoting the development of neurodegenerative diseases such as Alzheimer disease and PD.¹

We observed 3 new cases of parkinsonism among our cohort of 2,500 HIV-infected patients over a 2-year period, which is 4–8 times higher than the reported incidence of PD among people age 40–59 years.³ This estimate suggests HIV and HAART may preferentially predispose to the development of parkinsonism. Early HIV disease can rarely be associated with parkinsonism, probably reflecting viral involvement of the basal ganglia soon after seroconversion.⁴ Later in the course of HIV, in patients with HIV-associated dementia, features of parkinsonism may also occur, with postmortem studies confirming presence of HIV preferentially in inflammatory infiltrates and glial cells of the basal ganglia including the substantia nigra.⁵ The pattern of parkinsonism in the present series differs from these existing reports: 1) patients had chronic HIV infection with viral suppression in the plasma and minor replication in the CSF when symptoms developed, and 2) none had features of dementia. In fact, it remains possible that our patients have PD. We hypothesize that chronic HIV infection with very low level replication and HAART, particularly protease inhibitors, may predispose to neurodegeneration through persistent low level inflammation, mitochondrial dysfunction, and interference with the

ubiquitin proteasome pathway.¹ Proteasome dysfunction has been described at least with some of the protease inhibitors used in HAART. It may be advisable to avoid their use where possible in patients with parkinsonism.¹ In our patients, it was not possible to stop or reduce antiretroviral medications because of risk of HIV progression or the development of drug resistance. There is accumulating evidence that neurodegeneration occurs in patients with chronic HIV, even in the context of maximal viral suppression, at least in the plasma. Several groups have reported a 10%–40% incidence of cognitive impairment among virally suppressed chronically HIV-infected patients, particularly those with longer disease duration.⁶ Further evidence for neurodegeneration in patients with chronic HIV infection and HAART comes from a recent study which found that incidence of mild extrapyramidal signs is increased in patients with chronic HIV infection and HAART, particularly older patients and those with cognitive impairment.⁷ This can be interpreted as loss of nigrostriatal neuronal reserve in aging patients with HIV-HAART, which could predispose to the later development of parkinsonism. This view is supported by the significant finding of overexpression of neuritic alpha-synuclein in the substantia nigra of long surviving patients with HIV-HAART in the absence of HIV-associated neuropathology.² The present study extends these observations by confirming that chronic HIV infection with HAART may be associated with par-

kinsonism of relatively young onset, with rest tremor, that may be indistinguishable from idiopathic PD.

From the Department of Neurology, St Vincent's Hospital, University of New South Wales, Darlinghurst, Australia.

Disclosure: The authors report no disclosures.

Received January 18, 2009. Accepted in final form March 16, 2009.

Address correspondence and reprint requests to Dr. S. Tisch, Department of Neurology, St Vincent's Hospital, 390 Victoria Street, Darlinghurst, NSW 2010 Australia; stisch@stvincents.com.au

Copyright © 2009 by AAN Enterprises, Inc.

1. Brew BJ, Crowe SM, Landay A, Cysique LA, Guillemin G. Neurodegeneration and ageing in the HAART era. *J Neuroimmune Pharm* Epub 2008 Dec 6.
2. Khanlou N, Moore DJ, Chana G, et al. Increased frequency of alpha-synuclein in the substantia nigra in HIV infection. *J Neurovirol* 2009;15:131–138.
3. Van Den Eeden SK, Tanner CM, Bernstein AL, et al. Incidence of Parkinson's disease: variation by age, gender, and race/ethnicity. *Am J Epidemiol* 2003;157:1015–1022.
4. Hersh BP, Rajendran PR, Battinelli D. Parkinsonism as the presenting manifestation of HIV infection: improvement on HAART. *Neurology* 2001;56:278–279.
5. Brew BJ, Rosenblum M, Cronin K, Price RW. AIDS dementia complex and HIV-1 brain infection: clinical-virological correlations. *Ann Neurol* 1995;38:563–570.
6. Cysique LA, Maruff P, Brew J. Variable benefit in neuropsychological function in HIV-infected HAART-treated patients. *Neurology* 2006;66:1447–1450.
7. Valcour V, Watters MR, Williams AE, Sacktor N, McMurtry A, Shikuma C. Aging exacerbates extrapyramidal motor signs in the era of highly active antiretroviral therapy. *J Neurovirol* 2008;14:1–6.

Save These Dates for AAN CME Opportunities!

Mark these upcoming dates on your calendar for these exciting continuing education opportunities, where you can catch up on the latest neurology information.

AAN Regional Conference

- November 6–8, 2009, Las Vegas, Nevada, Planet Hollywood Resort and Casino

AAN Annual Meetings

- April 10–17, 2010, Toronto, Ontario, Canada, Toronto Convention Centre
- April 9–16, 2011, Honolulu, Hawaii, Hawaii Convention Center