

\*K Michael Cummings, Matthew J Carpenter  
 Department of Psychiatry and Behavioral Sciences (KMC, MJC) and  
 Cancer Control and Prevention, Hollings Cancer Center Medical  
 University of South Carolina (KMC, MJC), Medical University of  
 South Carolina, Charleston, SC 29425–8610, USA  
 cummingk@muscc.edu

KMC has received grant funding from Pfizer to study the effect of a hospital-based tobacco cessation intervention, and has received funding as an expert witness in litigation filed against the tobacco industry. MJC declares no competing interests.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND license.

- Hughes JR, Solomon LJ, Fingar JR, Naud S, Helzer JE, Callas PW. The natural history of efforts to stop smoking: a prospective cohort study. *Drug Alcohol Depend* 2013; **128**: 171–74.
- Kotz D, Brown J, West R. Prospective cohort study of the effectiveness of smoking cessation treatments used in the 'real world'. *Mayo Clin Proc* 2014; **89**: 1360–67.
- Dobbie F, Hiscock R, Leonardi-Bee J, et al. Evaluating long-term outcomes of NHS Stop Smoking Services (ELONS): a prospective cohort study. *Health Technol Assess* 2015; **19**: 1–156.
- West R, May S, West M, Croghan E, McEwen A. Performance of English stop smoking services in first 10 years: analysis of service monitoring data. *BMJ* 2013; **347**: f4921.
- Lifestyles Statistical team, Health and Social Care Information Centre. Statistics on smoking: England 2015. May 29, 2015. <http://content.digital.nhs.uk/catalogue/pub17526/stat-smok-eng-2015-rep.pdf> (accessed Nov 21, 2016).
- Gibson JE, Murray RL, Borland R, et al. The impact of the United Kingdom's national smoking cessation strategy on quit attempts and use of cessation services: findings from the International Tobacco Control Four Country Survey. *Nicotine Tob Res* 2010; **12** (suppl): S64–71.
- Hopkinson NS, Millett C, Glantz S, Arnott D, McNeill A. UK government should fund stop smoking media campaigns not give tax breaks to films with smoking imagery. *Addiction* 2016; **111**: 2066–67.
- Godfrey C, Parrott S, Coleman T, Pound E. The cost-effectiveness of the English smoking treatment services: evidence from practice. *Addiction* 2005; **100** (suppl 2): 70–83.
- Gilbert H, Sutton S, Morris R, et al. Effectiveness of personalised risk information and taster sessions to increase the uptake of smoking cessation services (Start2quit): a randomised controlled trial. *Lancet* 2017; published online Jan 24. [http://dx.doi.org/10.1016/S0140-6736\(16\)32379-0](http://dx.doi.org/10.1016/S0140-6736(16)32379-0).
- Tzelepis F, Paul CL, Walsh RA, McElduff P, Knight J. Proactive telephone counseling for smoking cessation: meta-analyses by recruitment channel and methodological quality. *J Natl Cancer Inst* 2011; **103**: 922–41.
- Fulton EA, Brown KE, Kwah KL, Wild S. StopApp: using the behaviour change wheel to develop an app to increase uptake and attendance at NHS Stop Smoking Services. *Healthcare* 2016; **4**: 31.
- Carpenter MJ, Hughes JR, Gray KM, Wahlquist AE, Saladin ME, Alberg AJ. Nicotine therapy sampling to induce quit attempts among smokers unmotivated to quit: a randomized clinical trial. *Arch Intern Med* 2011; **171**: 1901–07.
- Cancer Research UK, Action On Smoking And Health. Cutting down: the reality of budget cuts to local tobacco control. A survey of tobacco control leads in local authorities in England. November, 2016. [http://www.cancerresearchuk.org/sites/default/files/local\\_authority\\_survey\\_2016\\_report\\_cruk\\_finalfinal.pdf](http://www.cancerresearchuk.org/sites/default/files/local_authority_survey_2016_report_cruk_finalfinal.pdf) (accessed Nov 21, 2016).

## W Stressed brain, stressed heart?



pockygallery/123RF Stock Photo

In the past decade, more and more individuals experience psychosocial stress on a daily basis. Heavy workloads, job insecurity, or living in poverty are circumstances that can result in chronically increased stress, which in turn can lead to chronic psychological disorders such as depression. Besides the heavy psychological burden, chronic stress is also associated with an increased risk of cardiovascular disease.<sup>1,2</sup> Individuals with increased stress perception have a substantially higher prevalence of myocardial infarction than controls, as was reported by Rosengren and colleagues in 2004.<sup>2</sup> However, these association studies have thus far not established whether increased stress is causally related to the incidence of cardiovascular disease, and neither have they provided mechanistic insight into the pathways involved. This lack of mechanistic insight might be at least partly due to difficulties measuring personal stress levels—which can vary between individuals as stress perception differs from person to person—at the time of the stressor. Experimental work in animal models has established that both acute and chronic stress contribute to the underlying cause of cardiovascular disease<sup>3</sup>—ie, development of atherosclerosis, which is generally mediated by an

increased inflammatory response to a stressor. However, direct clinical evidence in human beings was lacking.

In *The Lancet*, Ahmed Tawakol and colleagues<sup>4</sup> report that the activity of the amygdala, a key component of the brain involved in emotion and stress, is of predictive value for the incidence of acute cardiovascular events. Their main study included 293 individuals without previous history of cardiovascular disease who underwent <sup>18</sup>F fluorodeoxyglucose (<sup>18</sup>F-FDG) PET or CT to visualise amygdalar activity as a measure for underlying stress. After median follow-up of 3.7 years, amygdalar activity was significantly associated with the incidence of cardiovascular events, including myocardial infarction, heart failure, and cerebrovascular events, even after multivariate adjustments. <sup>18</sup>F-FDG PET or CT also allows visualisation of haemopoietic tissue activity and the activity of inflammatory cells within the vessel wall. Statistical mediation analysis showed that bone-marrow activity, followed by vascular inflammation, was a significant mediator in the association between amygdalar activity and the incidence of cardiovascular events. Although these mediation analyses do not provide evidence of causality, the data are in line with

Published Online

January 11, 2017

[http://dx.doi.org/10.1016/S0140-6736\(17\)30044-2](http://dx.doi.org/10.1016/S0140-6736(17)30044-2)

See [Articles](#) page 834

data from an animal study,<sup>5</sup> in which stress induced an inflammatory response via activation of the bone marrow. Additionally, interleukin 6 derived from bone-marrow-derived leucocytes has been implicated as one of the causal mediators of the stress response in mice.<sup>6</sup> How the reported increase in the inflammatory response directly relates to the rise in acute cardiovascular syndromes caused by atherosclerotic plaque rupture or erosion<sup>7</sup> remains to be investigated.

Of clinical relevance is the perceived stress study that Takawol and colleagues also did,<sup>4</sup> which included patients with a history of post-traumatic stress disorder (PTSD). In these patients, perceived stress was associated with amygdalar activity, vascular inflammation, and circulating c-reactive protein (CRP) concentrations. The association between stress and CRP concentrations is in line with data obtained from other cohorts containing patients with PTSD, depression, and other diseases that are associated with chronic stress, in whom increased circulating CRP, interleukin 6, and tumour necrosis factor  $\alpha$  concentrations were noted.<sup>8-10</sup> The findings of these studies<sup>8-10</sup> suggest that individuals with high perceived stress levels—whether patients with PTSD, depression, or anxiety, or other individuals experiencing chronic stress—are at higher risk of having a cardiovascular disease event.

Although only 13 patients with PTSD were included in the perceived stress study (and no healthy controls with basal amygdalar activity), this study provides important evidence justifying the initiation of new studies of larger populations and with longer follow-up to assess cardiovascular disease risk directly in patients with PTSD. Because interleukin 6 is repeatedly increased in stress in both human beings and rodents, analysis of circulating interleukin 6 concentrations, next to CRP, might provide more insight into the underlying inflammatory mechanisms. In Takawol and colleagues' outcomes study,<sup>4</sup> the association between amygdalar activity and the incidence of cardiovascular disease still remains

after correction for history of depression or anxiety, which suggests that these data are not only applicable to patients with severe psychological disorders, but also to individuals without previous history of these conditions. Together, these clinical data establish a connection between stress and cardiovascular disease, thus identifying chronic stress as a true risk factor for acute cardiovascular syndromes, which could, given the increasing number of individuals with chronic stress, be included in risk assessments of cardiovascular disease in daily clinical practice.

*\*Ilze Bot, Johan Kuiper*

Division of Biopharmaceutics, Leiden Academic Centre for Drug Research, Leiden University, Einsteinweg 55, 2333 CC, Leiden, Netherlands  
i.bot@lacdr.leidenuniv.nl

We declare no competing interests.

- 1 Nabi H, Kivimaki M, Batty GD, et al. Increased risk of coronary heart disease among individuals reporting adverse impact of stress on their health: the Whitehall II prospective cohort study. *Eur Heart J* 2013; **34**: 2697-705.
- 2 Rosengren A, Hawken S, Ounpuu S, et al. Association of psychosocial risk factors with risk of acute myocardial infarction in 11 119 cases and 13 648 controls from 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; **364**: 953-62.
- 3 Lagraauw HM, Kuiper J, Bot I. Acute and chronic psychological stress as risk factors for cardiovascular disease: Insights gained from epidemiological, clinical and experimental studies. *Brain Behav Immun* 2015; **50**: 18-30.
- 4 Tawakol A, Ishai A, Takx RAP, et al. Relation between resting amygdalar activity and cardiovascular events: a longitudinal and cohort study. *Lancet* 2017; published online Jan 11. [http://dx.doi.org/10.1016/S0140-6736\(16\)31714-7](http://dx.doi.org/10.1016/S0140-6736(16)31714-7).
- 5 Heidt T, Sager HB, Courties G, et al. Chronic variable stress activates hematopoietic stem cells. *Nat Med* 2014; **20**: 754-58.
- 6 Hodes GE, Pfau ML, Leboeuf M, et al. Individual differences in the peripheral immune system promote resilience versus susceptibility to social stress. *Proc Natl Acad Sci U S A* 2014; **111**: 16136-41.
- 7 Hansson GK, Libby P, Tabas I. Inflammation and plaque vulnerability. *J Intern Med* 2015; **278**: 483-93.
- 8 Von Känel R, Hepp U, Kraemer B, et al. Evidence for low-grade systemic proinflammatory activity in patients with posttraumatic stress disorder. *J Psychiatr Res* 2007; **41**: 744-52.
- 9 Duivels HE, Vogelzangs N, Kupper N, de Jonge P, Penninx BW. Differential association of somatic and cognitive symptoms of depression and anxiety with inflammation: findings from the Netherlands Study of Depression and Anxiety (NESDA). *Psychoneuroendocrinology* 2013; **38**: 1573-85.
- 10 Fagundes CP, Glaser R, Hwang BS, Malarkey WB, Kiecolt-Glaser JK. Depressive symptoms enhance stress-induced inflammatory responses. *Brain Behav Immun* 2013; **31**: 172-76.

## Women's cancers: shining a light on a neglected health inequity

Breast and cervical cancers receive far less funding, advocacy, and public and political attention in low-income and middle-income countries (LMICs) than in high-income countries (HICs). Yet women in these settings have higher burdens of these diseases, poorer

access to care, present with more advanced stages of disease, and are more likely to die from their disease than women in HICs. Why the discrepancy? Reasons include a belief that breast and cervical cancers are not a priority when compared with other women's health

Published Online  
November 1, 2016  
[http://dx.doi.org/10.1016/S0140-6736\(16\)31798-6](http://dx.doi.org/10.1016/S0140-6736(16)31798-6)  
See [Comment](#) pages 773 and 775