

# Do people living with HIV experience greater age advancement than their HIV-negative counterparts?

Davide De Francesco<sup>a</sup>, Ferdinand W. Wit<sup>b,c</sup>, Alexander Bürkle<sup>d</sup>, Sebastian Oehlke<sup>d</sup>, Neeltje A. Kootstra<sup>e</sup>, Alan Winston<sup>f</sup>, Claudio Franceschi<sup>g</sup>, Paolo Garagnani<sup>g</sup>, Chiara Pirazzini<sup>g</sup>, Claude Libert<sup>h,i</sup>, Tilman Grune<sup>j</sup>, Daniela Weber<sup>j</sup>, Eugène H.J.M. Jansen<sup>k</sup>, Caroline A. Sabin<sup>a</sup>, Peter Reiss<sup>b,c</sup>,  
on behalf of the the Co-morBidity in Relation to AIDS (COBRA) Collaboration

See related paper on page 345

**Objectives:** Despite successful antiretroviral therapy, people living with HIV (PLWH) may show signs of premature/accelerated aging. We compared established biomarkers of aging in PLWH, appropriately chosen HIV-negative individuals, and blood donors, and explored factors associated with biological age advancement.

**Design:** Cross-sectional analysis of 134 PLWH on suppressive antiretroviral therapy, 79 lifestyle-comparable HIV-negative controls aged 45 years or older from the Co-morBidity in Relation to AIDS (COBRA) cohort, and 35 age-matched blood donors.

**Methods:** Biological age was estimated using a validated algorithm based on 10 biomarkers. Associations between 'age advancement' (biological minus chronological age) and HIV status/parameters, lifestyle, cytomegalovirus (CMV), hepatitis B (HBV) and hepatitis C virus (HCV) infections were investigated using linear regression.

**Results:** The average (95% CI) age advancement was greater in both HIV-positive [13.2 (11.6–14.9) years] and HIV-negative [5.5 (3.8–7.2) years] COBRA participants compared with blood donors [−7.0 (−4.1 to −9.9) years, both  $P$ 's < 0.001], but also in HIV-positive compared with HIV-negative participants ( $P$  < 0.001). Chronic HBV, higher anti-CMV IgG titer and CD8<sup>+</sup> T-cell count were each associated with increased age advancement, independently of HIV-status/group. Among HIV-positive participants, age advancement was increased by 3.5 (0.1–6.8) years among those with nadir CD4<sup>+</sup> T-cell count less than 200 cells/ $\mu$ l and by 0.1 (0.06–0.2) years for each additional month of exposure to saquinavir.

<sup>a</sup>Institute for Global Health, University College London, London, UK, <sup>b</sup>Department of Global Health, Academic Medical Center and Amsterdam Institute for Global Health and Development, <sup>c</sup>Stichting HIV Monitoring, Amsterdam, The Netherlands, <sup>d</sup>Molecular Toxicology Group, University of Konstanz, Konstanz, Germany, <sup>e</sup>Department of Experimental Immunology, Academic Medical Center, Amsterdam, The Netherlands, <sup>f</sup>Division of Infectious Diseases, Imperial College London, London, UK, <sup>g</sup>Department of Experimental, Diagnostic and Specialty Medicine, Alma Mater Studiorum Università di Bologna, Bologna, Italy, <sup>h</sup>Department of Biomedical Molecular Biology, Ghent University, <sup>i</sup>Center for Inflammation Research, Flanders Institute for Biotechnology, Ghent, Belgium, <sup>j</sup>Department of Molecular Toxicology, German Institute of Human Nutrition, Nuthetal, Germany, and <sup>k</sup>National Institute for Public Health and the Environment, Bilthoven, The Netherlands.

Correspondence to Davide De Francesco, Institute for Global Health, University College London, Royal Free Campus, Rowland Hill Street, London NW3 2PF, UK.

Tel: +44 20 7794 0500 x38827; e-mail: d.defrancesco@ucl.ac.uk

Received: 3 May 2018; accepted: 28 June 2018.

DOI:10.1097/QAD.0000000000002063

ISSN 0269-9370 Copyright © 2018 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

**Conclusion:** Both treated PLWH and lifestyle-comparable HIV-negative individuals show signs of age advancement compared with blood donors, to which persistent CMV, HBV co-infection and CD8<sup>+</sup> T-cell activation may have contributed. Age advancement remained greatest in PLWH and was related to prior immunodeficiency and cumulative saquinavir exposure. Copyright © 2018 The Author(s). Published by Wolters Kluwer Health, Inc.

*AIDS* 2019, **33**:259–268

**Keywords:** accelerated aging, aging, biological age, biomarkers of aging, HIV, premature aging

## Introduction

Despite the success of combination antiretroviral therapy, people living with HIV (PLWH) have an increased burden of noncommunicable age-associated comorbidities compared with HIV-negative individuals [1,2]. The causes of this increased burden of comorbidities remain unclear but may involve an accelerated or accentuated aging process [3,4], resulting from a complex mix of HIV infection, antiretroviral treatment, chronic viral co-infections and lifestyle/behavioral factors.

Aging can be defined as the time-dependent decline of functional capacity and stress resistance associated with increased risk of disability, morbidity and mortality [5]. There is clear evidence that the rate of aging differs significantly between individuals, because of genetic heterogeneity and environmental factors [6]. Therefore, chronological age may not represent the best way of measuring aging and may not accurately reflect an individual's position in his/her total lifespan [7]. This has led to a search for reliable biomarkers of aging, defined as biological parameters that capture the age-related changes in body function or composition. These biomarkers could serve to measure 'biological' age, a hypothetical value denoting the extent of age-related changes in function and composition of a human body, and predict the onset of age-related diseases and/or expectant residual lifetime more accurately than chronological age [8].

Many candidate biomarkers of aging have been proposed in the scientific literature and have been used to investigate the association between HIV and aging [9]. Among these, there are markers of chronic systemic immune activation (soluble CD14<sup>+</sup> and CD163<sup>+</sup> [10,11]), inflammation (C-reactive protein and interleukin-6 [12]), coagulation (D-dimer [13]), leukocyte telomere length [14], somatic mitochondrial DNA mutations [15], expression levels of the cell cycle regulator CDKN2A [14], DNA methylation levels [16], ophthalmological parameters [17] and age-related brain atrophy [18].

Combinations of biomarkers may more reliably measure these age-related changes (or what is called also 'biological age'), giving more accurate estimates of residual lifetime than that obtained from any single

biomarker in isolation. The MARK-AGE project has proposed a method to combine powerful biomarkers of human aging (most of which have been shown to be good markers of age when taken in isolation [19,20]) to predict biological age of individuals, and therefore, assess the aging process [21,22].

The current study aimed to compare established biomarkers of aging and their combination (as proposed by the MARK-AGE study) in order to evaluate the biological age of PLWH, demographically and lifestyle-matched HIV-negative individuals, and blood donors with similar chronological age. Furthermore, we investigated the associations between any observed age advancement and lifestyle risk factors, chronic viral co-infections including hepatitis B virus (HBV), hepatitis C virus (HCV) and cytomegalovirus (CMV), HIV-related parameters and (cumulative) past or current exposure to antiretroviral drugs.

## Methods

### Study participants

HIV-positive ( $n = 134$ ) and HIV-negative individuals with similar sociodemographic and lifestyle factors ( $n = 79$ ) were prospectively enrolled in the COmorBidity in Relation to AIDS (COBRA) cohort from HIV outpatient clinics and sexual health centres, respectively, located in Amsterdam (Netherlands) and London (UK) [23]. Inclusion criteria were age 45 years or older ( $\geq 50$  years in London), laboratory-confirmed presence or absence of HIV-1 infection, and the HIV-positive people were required to have plasma HIV-RNA less than 50 copies/ml for at least 12 months on antiretroviral therapy. Exclusion criteria were: current major depression [as indicated by a Patient Health Questionnaire-9 (PHQ-9) score  $\geq 15$  at screening], confounding neurological diseases, previous severe head injury (with loss of consciousness for  $\geq 30$  min), a history of cerebral infections (including AIDS-defining diseases involving the central nervous system), current intravenous drug use (in the past 6 months, based on self-report), daily use of recreational drugs (with the exception of cannabis), excess alcohol intake ( $>48$  units per week), severe

psychiatric disease or contraindication to MRI scan or lumbar puncture examination. Recruitment took place between December 2011 and October 2014. The study was approved by the institutional review board of the Academic Medical Center (AMC) in Amsterdam (reference number NL 30802.018.09) and a UK Research Ethics Committee (REC) (reference number 13/LO/0584 Stanmore, London). All participants gave written informed consent.

In addition, 35 blood donors were selected from the Dutch national blood bank in Amsterdam, the Netherlands (<https://www.sanquin.nl/en/>). Blood donors were age-matched with the HIV-positive and HIV-negative COBRA participants and, as a requirement for blood donation in the Netherlands, they had screened negative for HIV, HBV, HCV, syphilis, and human T-lymphotropic virus 1 and 2 (HTLV) infections. Candidate blood donors in the Netherlands are also eventually excluded from blood donation based on a questionnaire with regard to health (general and sexual), medication use, sexual risk behavior and travel.

### Biomarkers of aging

A 10-item panel of biomarkers of aging identified by the MARK-AGE project [21,22] were measured and are listed in Supplementary Table 1, <http://links.lww.com/QAD/B389>. Briefly, these biomarkers have been selected as best predictors of chronological age among nearly 400 candidate biomarkers in a population of approximately 3300 individuals aged between 30 and 74 years (mean age was 56 years) recruited from eight European countries. Lycopene, and  $\alpha$ -tocopherol in plasma were simultaneously determined by high performance liquid chromatography with UV and fluorescence detection as previously described [24]. Alpha-2-macroglobulin in plasma was measured on an autoanalyzer (DxC 800; Beckman-Coulter, Woerden, The Netherlands) by an immunoturbidimetric method using reagents from Dialab, Wiener Neudorf, Austria as described by Jansen *et al.* [25]. Dehydroepiandrosterone sulfate, ferritin (women only), and prostate-specific antigen (men only) in plasma were analyzed using an immuno-analyzer (Access-2; Beckman-Coulter) [25]. ELOVL2 and FHL2 DNA methylation in purified PBMC were analyzed using the Agena Bioscience's EpiTYPER DNA methylation analysis technology [26]. The N-glycans present on the proteins in plasma were released, labelled, and analyzed by DSA-FACE technology, as described previously [20].

### Age advancement

The biological age of each individual was derived separately for men and women as a linear combination of these biomarkers using the method and the weights described in Bürkle *et al.* [22]. Age advancement for each individual was then defined as the difference between biological and chronological age. Positive age

advancement is, therefore, indicative of more age-related changes in body function and composition than what would be expected in the average population, given the observed chronological age; negative age advancement would suggest less age-related changes.

### Statistical analysis

Participants' characteristics and distribution of the 10 biomarkers used to derive biological age are reported as raw and relative frequencies or median and interquartile range (IQR) and differences across groups were evaluated using chi-square or Wilcoxon signed-rank tests, as appropriate. Age advancement in each group is reported as mean and 95% confidence interval (CI) and assessed for significance using the one-sample *t*-test. Differences between groups were assessed using linear regression with HIV status/study group as categorical independent variable.

Associations were considered between age advancement and the following factors:

- (1) Sociodemographic factors (age, sex, ethnicity, sexuality, years of education)
- (2) Self-reported smoking, alcohol consumption and current recreational drug use
- (3) CMV (CMV serostatus, CMV total IgG and CMV high-avidity IgG titers), chronic HBV (defined as detectable hepatitis B surface antigen and/or HBV DNA) and HCV (defined as detectable HCV RNA)
- (4) HIV disease parameters such as CD4<sup>+</sup> and CD8<sup>+</sup> T-cell counts, CD4<sup>+</sup> : CD8<sup>+</sup> T-cell ratio, nadir CD4<sup>+</sup> T-cell count, nadir CD4<sup>+</sup> T-cell count <200 cells/ $\mu$ l, years since HIV diagnosis, prior AIDS diagnosis (defined as prior category C event as per the Centers for Disease Control and Prevention's classification system for HIV infection)
- (5) Exposure to antiretroviral drugs considered as both prior exposure (yes or no) and months of cumulative exposure.

The associations of age advancement with each factor, unadjusted and adjusted for HIV status/study group, were assessed through a series of linear regression models (with each factor considered one at the time in separate models). Factors that remained significantly associated with age advancement after adjustment for HIV status and group ( $P < 0.1$ ), were then included in a multivariable regression model to identify factors independently associated with age advancement. All statistical analyses were performed using SAS v9.4 (Cary, North Carolina, USA).

## Results

### Cohort characteristics

The 79 HIV-negative COBRA participants were comparable with the 134 HIV-positive participants in terms of age, sex, years of education, smoking and recreational drug use. HIV-positive participants were

**Table 1. Characteristics of the COBRA participants and blood donors ( $P^1$ : HIV-positive vs. HIV-negative COBRA participants;  $P^2$ : HIV-positive COBRA participants vs. blood donors: both were obtained from chi-square or Wilcoxon signed-rank tests, as appropriate).**

| <i>n</i> (%) or median (IQR)                                     | COBRA participants             |                               | $P^1$  | Blood donors ( <i>n</i> = 35) | $P^2$  |
|--|--------------------------------|-------------------------------|--------|-------------------------------|--------|
|  | HIV-positive ( <i>n</i> = 134) | HIV-negative ( <i>n</i> = 79) |        |                               |        |
| Age (years)  | 55 (51–62)                     | 57 (52, 64)                   | 0.24   | 59 (52, 65)                   | 0.37   |
| Sex: female  | 9 (6.7%)                       | 6 (7.6%)                      | 0.79   | 17 (48.6%)                    | <0.001 |
| Male   | 125 (93.3%)                    | 73 (92.4%)                    |        | 18 (51.4%)                    |        |
| Ethnicity: black-African   | 16 (12.0%)                     | 2 (2.6%)                      | 0.03   | n/a                           |        |
| White  | 117 (88.0%)                    | 76 (97.4%)                    |        | n/a                           |        |
| Sexuality: MSM   | 104 (77.6%)                    | 59 (74.7%)                    | 0.45   | n/a                           |        |
| Bisexual   | 10 (7.5%)                      | 4 (5.1%)                      |        | n/a                           |        |
| Heterosexual   | 18 (13.4%)                     | 16 (20.2%)                    |        | n/a                           |        |
| Years of education   | 14 (13, 16)                    | 16 (14, 17)                   | 0.23   | n/a                           |        |
| Smoking status: current smoker                                   | 40 (29.9%)                     | 20 (25.3%)                    | 0.24   | n/a                           |        |
| Ex-smoker  | 58 (43.2%)                     | 29 (36.7%)                    |        | n/a                           |        |
| Never smoked   | 36 (26.9%)                     | 30 (38.0%)                    |        | n/a                           |        |
| Alcohol consumption: current drinker                             | 104 (77.6%)                    | 71 (89.9%)                    | 0.04   | n/a                           |        |
| Previous drinker   | 18 (13.4%)                     | 3 (3.8%)                      |        | n/a                           |        |
| Never drunk  | 12 (9.0%)                      | 4 (5.1%)                      |        | n/a                           |        |
| Recreational drugs use in past 6 months                          | 44 (32.8%)                     | 18 (22.8%)                    | 0.16   | n/a                           |        |
| Chronic HBV infection  | 7 (5.3%)                       | 0 (0.0%)                      | 0.05   | 0 (0.0%)                      | 0.34   |
| Chronic HCV infection  | 5 (3.7%)                       | 0 (0.0%)                      | 0.16   | 0 (0.0%)                      | 0.58   |
| CMV infection  | 129 (97.7%)                    | 63 (79.8%)                    | <0.001 | 8 (22.9%)                     | <0.001 |
| Total anti-CMV IgG (AU)  | 51.2 (29.0–107.0)              | 21.1 (11.8–53.8)              | <0.001 | 11.3 (10.2–16.8)              | 0.003  |
| High avidity anti-CMV IgG (AU)                                   | 29.4 (14.9–61.1)               | 12.6 (7.2–25.3)               | <0.001 | 10.7 (10.0–13.2)              | 0.02   |
| CD4 <sup>+</sup> T-cell count (cells/ $\mu$ l)                   | 618 (472–806)                  | 900 (692–1174)                | <0.001 | n/a                           |        |
| CD8 <sup>+</sup> T-cell count (cells/ $\mu$ l)                   | 770 (611– 947)                 | 422 (313–614)                 | <0.001 | n/a                           |        |
| CD4 <sup>+</sup> : CD8 <sup>+</sup> T-cell count ratio           | 0.84 (0.60–1.12)               | 2.01 (1.44–2.64)              | <0.001 | n/a                           |        |
| Nadir CD4 <sup>+</sup> T-cell count (cells/ $\mu$ l)             | 180 (90–250)                   | n/a                           |        | n/a                           |        |
| Years since HIV diagnosis  | 15.0 (9.1–20.0)                | n/a                           |        | n/a                           |        |
| Duration of antiretroviral therapy (years)                       | 12.5 (7.4–16.9)                | n/a                           |        | n/a                           |        |
| Nadir CD4 <sup>+</sup> T-cell count less than 200 cells/ $\mu$ l | 83 (61.9%)                     | n/a                           |        | n/a                           |        |
| Prior AIDS   | 42 (31.3%)                     | n/a                           |        | n/a                           |        |

ARV, antiretroviral therapy; COBRA, Co-morBidity in Relation to AIDS; CMV, cytomegalovirus; HBV, hepatitis B virus; HCV, hepatitis C virus.

more likely to be of black-African origin ( $P=0.03$ ) and were less likely to report current alcohol consumption ( $P=0.04$ ) than the HIV-negative participants. Compared with both groups of COBRA participants, the blood donor group included a greater proportion of women ( $P<0.001$ ). CMV, chronic HBV and HCV co-infections were all more frequent in HIV-positive COBRA participants compared with both the HIV-negative participants and blood donors, with CMV also being more frequent in HIV-negative participants than in blood donors. All HIV-positive COBRA participants had plasma HIV RNA less than 50 copies/ml and were on antiretroviral therapy at study visit, they had a median (IQR) CD4<sup>+</sup> T-cell count of 618 (472–806) cells/ $\mu$ l, and 31% had a prior clinical AIDS diagnosis (Table 1). The number and proportion of people who had ever been exposed to each antiretroviral drug and the median (IQR) duration of exposure are reported in Supplementary Table 2, <http://links.lww.com/QAD/B389>.


### Biomarkers of aging

The distribution of each of the 10 biomarkers of aging used to derive biological age is summarized in Supplementary Table 3, <http://links.lww.com/QAD/B389>. HIV-positive COBRA participants had a greater cumulative proportion of cytosine methylation at four out

of five gene positions compared with HIV-negative COBRA participants (all  $P$ 's  $\leq 0.01$ ). DHEAS and A2M (in male participants only) concentrations were also significantly different in the two COBRA groups ( $P=0.03$  and  $P=0.004$ , respectively). In all the five gene positions, the cumulative proportion of cytosine methylation in blood donors was significantly lower than that seen in both COBRA groups. A2M concentration (in males only) in blood donors was lower compared with both COBRA groups (both  $P$ 's  $<0.001$ ), whereas N-glycan peak 6 was higher compared with COBRA HIV-positive ( $P=0.04$ ) but not with HIV-negative participants ( $P=0.18$ ).

### Age advancement in HIV-positive and HIV-negative COBRA participants and blood donors

Biological age was significantly greater than chronological age by a mean of 13.2 (95% CI 11.6–14.9) years in HIV-positive COBRA participants and by 5.5 (3.8–7.2) years in HIV-negative participants ( $P<0.001$  for each). In contrast, biological age was a mean of 7.0 (4.1–9.9) years lower than chronological age in blood donors ( $P<0.001$ , Fig. 1a). Whilst age advancement was greater in both COBRA groups compared with blood donors ( $P<0.001$  for each), the HIV-positive COBRA



**Fig. 1.** (a) Age advancement (biological minus chronological age) in HIV-positive and HIV-negative COBRA participants and blood donors ( $P$ s from linear regression); (b) Correlation between age advancement and chronological age in HIV-positive and HIV-negative COBRA participants and blood donors (no interaction between chronological age and HIV-status/group,  $P = 0.66$ ). BD, blood donors; COBRA, Co-morBidity in Relation to AIDS.

participants also demonstrated greater age advancement than the HIV-negative participants ( $P < 0.001$ ). Age advancement was also negatively correlated with chronological age (Pearson's  $r = -0.17$ ,  $P = 0.08$ ) with no significant interaction with HIV-status/group ( $P = 0.66$ , Fig. 1b).

### Factors associated with age advancement

Across the entire study population, no significant associations were found between age advancement and ethnicity, sexual orientation or lifestyle factors (Table 2). Overall, men had greater age advancement than women even after adjustment for HIV-status/group. However, among COBRA participants only, the difference (95% CI) between men and women was only 0.09 (−5.04 to 5.22) years ( $P = 0.97$ ) and the difference between COBRA groups and blood donors remained significant even after adjusting for sex (both  $P$ s  $< 0.001$ ). Viral co-infections such as CMV and chronic HBV, as well as  $CD4^+$  and  $CD8^+$  T-cell count and their ratio appeared to be associated with increased age advancement. However, only chronic HBV, total and high avidity anti-CMV IgG antibody titer and  $CD8^+$  T-cell count showed significant associations, which were also independent of HIV-status/group.

In multivariable analysis including significant factors after adjusting for HIV-status/group ( $P < 0.1$ ), the average increase in age advancement (95% CI) remained significant for chronic HBV [10.05 (3.45–16.64) years,  $P = 0.003$ ], total anti-CMV IgG antibody titer [1.83 (0.51–3.15) years per one log (AU) increase,  $P = 0.007$ ] and  $CD8^+$  T cell [0.44 (0.02–0.85) years per 100 cells/ $\mu$ l increase,  $P = 0.04$ ]. Even after adjusting for chronic HBV,

total anti-CMV IgG and  $CD8^+$  T-cell, HIV-positive COBRA participants exhibited a greater age advancement compared with both the HIV-negative participants [mean increase (95% CI) 4.5 (1.6–7.5),  $P = 0.003$ ] and blood donors [mean increase (95% CI): 19.0 (12.6–25.4),  $P < 0.001$ ], with the HIV-negative COBRA participants also exhibiting greater age advancement than blood donors [mean increase (95% CI) 13.5 (7.1–20.0),  $P < 0.001$ ].

### Associations with HIV parameters and exposure to antiretroviral drugs

Among the HIV-positive participants in COBRA, in univariable regression analyses, positive correlations were found between age advancement and time since HIV diagnosis, duration of antiretroviral therapy and nadir  $CD4^+$  T-cell count  $< 200$  cells/ $\mu$ l (Table 3). Cumulative exposure and/or prior exposure to some antiretroviral drugs was also associated with an increased age advancement (Supplementary Table 4, <http://links.lww.com/QAD/B389>); in particular each additional year of exposure to saquinavir was associated with a 1.39 (95% CI 0.71–2.07) years increase in age advancement ( $P < 0.001$ ), but also prior exposure to stavudine or any d-drug were associated with greater age advancement ( $P = 0.02$  and  $P = 0.03$ , respectively). Due to the strong correlations between these factors, we ran a multivariable regression analysis with all HIV-parameters and exposures to antiretroviral drugs that were significantly associated with age advancement in univariable analysis ( $P < 0.1$ ), also accounting for chronic HBV, total anti-CMV IgG antibody titer and  $CD8^+$  T-cell count. In this multivariable analysis, only cumulative exposure to saquinavir (average increase of 1.17 (0.49, 1.85) years per year of

**Table 2. Mean (95% confidence interval) increase (if positive) or decrease (if negative) in age advancement among HIV-positive and HIV-negative COBRA participants and blood donors (combined), unadjusted, and adjusted for HIV status/group.**

| Variable  | Mean (95% CI) increase/decrease (years) |          |                               |          |
|---|---|----------|-------------------------------|----------|
|   | Unadjusted                              | <i>P</i> | Adjusted for HIV status/group | <i>P</i> |
| Age (per 5 years)   | -0.17 (-0.35 to 0.02)                   | 0.08     | -0.11 (-0.26 to 0.04)         | 0.15     |
| Gender: male vs. female   | 4.19 (0.01-8.36)                        | 0.05     | 4.55 (0.92-8.18)              | 0.01     |
| Ethnicity: black-African vs. white                                  | 3.40 (-1.28 to 8.09)                    | 0.15     | 1.18 (-3.21 to 5.56)          | 0.60     |
| Sexuality: MSM vs. non-MSM  | 0.92 (-1.86 to 3.69)                    | 0.51     | -0.01 (-3.23 to 3.20)         | 0.99     |
| Smoking: current smoker vs. never smoked                            | 2.46 (-0.95 to 5.88)                    | 0.16     | 1.47 (-1.69 to 4.64)          | 0.36     |
| Ex-smoker vs. never smoked  | 0.71 (-2.41 to 3.83)                    | 0.65     | -0.28 (-3.18 to 2.62)         | 0.85     |
| Alcohol: current drinker vs. never drunk                            | -1.80 (-6.81 to 3.21)                   | 0.48     | -0.58 (-5.24 to 4.08)         | 0.81     |
| Previous drinker vs. never drunk                                    | 0.60 (-5.57 to 7.16)                    | 0.81     | -0.03 (-5.93 to 8.87)         | 0.99     |
| Recreational drugs use  | -0.88 (-3.77 to 2.01)                   | 0.55     | -1.80 (-4.48 to 0.87)         | 0.18     |
| Chronic HBV infection   | 14.47 (6.12- 22.83)                     | <0.001   | 9.11 (2.41-15.82)             | 0.008    |
| Chronic HCV infection   | 6.50 (-3.50-16.50)                      | 0.20     | 1.09 (-6.89 to 9.08)          | 0.79     |
| CMV infection   | 11.46 (8.18-14.73)                      | <0.001   | 0.54 (-3.11 to 4.19)          | 0.77     |
| Total anti-CMV IgG [per 1 log (AU)]                                 | 3.49 (2.15-4.85)                        | <0.001   | 2.05 (0.73-3.36)              | 0.002    |
| High avidity anti-CMV IgG (per 1 AU)                                | 0.06 (0.03-0.09)                        | <0.001   | 0.04 (0.01-0.07)              | 0.01     |
| CD4 <sup>+</sup> T-cell count (per 100 cells/ $\mu$ l)              | -0.85 (-1.30 to -0.41)                  | <0.001   | -0.31 (-0.79 to 0.17)         | 0.21     |
| CD8 <sup>+</sup> T-cell count (per 100 cells/ $\mu$ l)              | 0.91 (0.53-1.28)                        | <0.001   | 0.49 (0.07-0.90)              | 0.02     |
| CD4 <sup>+</sup> : CD8 <sup>+</sup> T-cell count ratio (per 1-unit) | -2.87 (-4.08 to -1.66)                  | <0.001   | -1.11 (-2.59 to 0.38)         | 0.14     |

Blood donors were included only in analyses concerning age, sex, chronic hepatitis B virus (HBV) and hepatitis C virus (HCV) infection, cytomegalovirus (CMV) infection and anti-CMV IgG. Estimates were obtained with separate linear regression models (one for each factor considered).

exposure,  $P < 0.001$ ) and nadir CD4<sup>+</sup> T-cell count less than 200 cells/ $\mu$ l (average increase of 3.00 (-0.22 to 6.22) years,  $P = 0.07$ ), as well as chronic HBV (7.35 (0.42-14.29) years,  $P = 0.04$ ) and total anti-CMV IgG antibody titer (1.86 (0.22-3.51) years per one log increase,  $P = 0.03$ ), retained their significant associations.

## Discussion

Both successfully treated PLWH and people without HIV with similar lifestyles show signs of age advancement compared with healthy blood donors who, conversely, appear younger than their chronological age. This may be explained by the strict requirements around blood donation in the Netherlands. This hypothesis is supported, for example, by the observation that the prevalence of CMV infection was higher in HIV-negative participants (79.8%), and lower in blood donors (22.9%), than in the general Dutch population (50% prevalence at 50 years of age) [27].

Whilst both groups of COBRA participants exhibited age advancement, this was significantly greater in HIV-positive than in HIV-negative participants. Whilst this difference in age advancement did not appear to be explained by differences in participant characteristics, chronic viral co-infections such as CMV and HBV, prior immunosuppression and exposure to some antiretroviral drugs may have contributed to this age advancement. CMV and HBV co-infections may cause premature aging of the immune system in PLWH by their chronic antigenic stimulation, inducing systemic immune activation [28]. In particular, CMV reactivation and concurrent immune responses to control infection are associated with aging and increased morbidity and mortality in both the general population and PLWH [29,30]. In the same group of people reported here, we previously found that increased anti-CMV IgG levels are associated with a higher proportion of terminally differentiated CD4<sup>+</sup> and CD8<sup>+</sup> T cells as well as CD4<sup>+</sup> T-cell activation [31], which may, in turn, have contributed to the observed greater age advancement.

**Table 3. Mean (95% confidence interval) increase (if positive) or decrease (if negative) in age advancement among HIV-positive COBRA participants associated with each HIV-related factor.**

| Variable   | Mean (95% CI) increase/decrease (in years) | <i>P</i> |
|--|--|----------|
| Years since HIV diagnosis (per year)                             | 0.28 (-0.01-0.56)                          | 0.05     |
| Duration of antiretroviral therapy (per year)                    | 0.25 (0.01-0.49)                           | 0.05     |
| Nadir CD4 <sup>+</sup> T-cell count (per 100 cells/ $\mu$ l)     | -0.73 (-1.87-0.41)                         | 0.21     |
| Nadir CD4 <sup>+</sup> T-cell count less than 200 cells/ $\mu$ l | 3.48 (0.14-6.83)                           | 0.04     |
| Prior AIDS   | 1.70 (-1.85-5.24)                          | 0.35     |

Estimates were obtained with separate linear regression models (one for each factor considered). CI, confidence interval; COBRA, Co-morbidity in Relation to AIDS.

Interestingly, current or past smoking did not appear to affect age advancement. Whilst more direct and accurate measures of smoking (including duration and frequency of smoking) would be more appropriate to evaluate the effect on the aging process, we cannot exclude that our findings reflect a lack of association with markers used in this study, especially in cohorts with relatively low smoking frequency (COBRA participants who self-reported current smoking, reported a mean of 12 cigarettes smoked per day). Other factors that have been shown to be associated with different aspects of the aging process of PLWH in previous studies (i.e. HCV co-infection [32] and  $CD4^+$  :  $CD8^+$  T-cell count ratio [33]) were not linked to greater age advancement. Whilst this is likely to be because of a combination of low statistical power (only five participants were infected with chronic HCV) and collinearity with other lifestyle factors (e.g. recreational drugs for HCV and  $CD8^+$  T-cell count for  $CD4^+$  :  $CD8^+$  ratio), further studies are needed to elucidate potential contributors to the age advancement seen in PLWH.

Among antiretroviral drugs, we found a significant association between age advancement and duration of past exposure to saquinavir, which was independent of concomitant exposure to other drugs generally considered to have the greatest mitochondrial toxicities and potential effects on aging such as didanosine, stavudine, zalcitabine and zidovudine (data not shown). Whereas saquinavir has been shown to directly induce vascular endothelial toxicity *in vitro* [34], other HIV protease inhibitors were shown to be able to induce vascular smooth muscle cell senescence by downregulating ZMPSTE24, leading to prelamin A accumulation and potential premature vascular aging, changes, which were also observed in peripheral blood mononuclear cells from HIV-infected patients who were treated with the same protease inhibitors [35]. The reason why we only observed an association between age advancement and exposure to saquinavir, but not other HIV protease inhibitors, remains unclear, and suggests this observation should be interpreted with caution. Importantly, having experienced more pronounced immunodeficiency in HIV-positive participants was also linked to an increased age advancement suggesting that a greater effect on aging is likely to occur as the infection remains untreated for longer.

Our findings are consistent with previous reports of brain aging in the COBRA study [18] and with other studies of cellular and molecular markers of biological aging. Indeed, studies of telomere length and CDKN2A [14,36],  $CD8^+$  T-cell senescence [37], and DNA methylation profiles ('epigenetic clock') [16,38] showed similar indications of age advancement in PLWH. Other studies reported evidence of accentuated aging only in PLWH with low nadir  $CD4^+$  T-cell counts compared with HIV-negative individuals [39,40]. Of note, however, these

previous studies often included untreated PLWH (some with detectable HIV RNA) and HIV-negative controls that differed with regards to some lifestyle behaviors, such as smoking and alcohol consumption. In our study, effectively treated PLWH with high  $CD4^+$  T-cell counts and highly comparable HIV-negative controls were purposely recruited in order to reduce the influence of sociodemographic and lifestyle confounding factors. The importance of an appropriately chosen control group of HIV-negative individuals is also highlighted by the finding that HIV-negative COBRA participants showed greater age advancement compared with blood donors. Furthermore, these studies only focus on one single tissue or body system, and are therefore unable to reflect the intrinsic multicausal and multisystem nature of the aging process [41]. Chronic HIV may differently affect the rate of aging at the levels of cells, tissues or body systems within the same organism and this complexity is more likely reflected by a method used in our study, which integrates multiple sources of molecular, cellular and physiologic data.

Age advancement reflects the difference in body function and composition of an individual relative to that of a similarly-aged 'healthy' individual. As such, it is not surprising that we found a weak negative correlation with chronological age that, if anything, may reflect survivorship bias. Moreover, as we found no interaction between the correlation of age advancement with chronological age and HIV-status/group, our results are more suggestive of accentuated rather than accelerated aging in the context of treated HIV disease. However, longitudinal follow-up is required to more appropriately address this issue. Accentuated aging occurs when there is an increased burden of aging-related damage but the year-on-year damage remains static over time whereas accelerated aging occurs when the decline arises earlier than expected and implies a progressive increase in the rate of decline [42].

Unmeasured confounders (e.g. living settings (urban vs. rural), diet, physical activity, sleep habits and direct, rather than self-reported, information on smoking and substance/alcohol use) may also have contributed to the increased age advancement we observed in both PLWH and lifestyle-matched HIV-negative individuals and also to the apparent increase due to HIV. Our study has further limitations. Given the cross-sectional design, it can only assess associations and longitudinal studies would be necessary to evaluate the potential causal role of HIV and the interplay with antiretroviral drugs and viral co-infections. The sample size was believed to be sufficient to allow meaningful analysis; however, the study may not be powered enough to evaluate associations with cumulative exposure to antiretroviral drugs as they were evaluated only on participants with prior exposure to a drug. Also our cohort predominantly constitutes white, northern European MSM over the age of 45 years. Results may,

therefore, not be generalizable to younger populations or populations within a different HIV epidemic setting. Also, individuals with major depression were excluded from the study; as psychological stress has been linked with molecular changes that can affect the aging process [43], this could have resulted in an underestimation of age advancement in both COBRA groups. Finally, the age advancement observed in PLWH, COBRA HIV-negative individuals and blood donors is relative to the population used by the MARK-AGE project to develop the algorithm to estimate biological age. Individuals with self-reported HIV, HBV (except seropositivity by vaccination), HCV or cancer were excluded but the representativeness of the sample to European countries involved in the MARK-AGE study still needs to be assessed. Whilst the algorithm has not been yet validated in external population (including cohort of PLWH), it showed promising results in individuals affected by Down's syndrome who, as expected, showed greater age advancement compared with the general population (manuscript in preparation). Moreover, sensitivity analysis in our cohort suggest a positive correlation between age advancement and the number of age-associated comorbidities (Pearson's  $r=0.18$ ,  $P=0.007$ ) and time needed to walk 15 feet (Pearson's  $r=0.13$ ,  $P=0.04$ ) as well as a negative correlation with hand grip (Pearson's  $r=-0.12$ ,  $P=0.05$ ), a validated measure of frailty.

In conclusion, our results suggest that PLWH with undetectable plasma HIV RNA may experience accentuated aging compared with HIV-negative individuals with similar lifestyles, as estimated using a set of validated biomarkers of aging. This age advancement appears to be related to viral co-infections such as CMV and chronic HBV, but also to historic severe immunosuppression and possibly exposure to particular antiretroviral drugs. Future longitudinal studies are required to further help clarifying the effect of HIV and its treatment on the natural aging process and the functional and clinical consequences in the millions of PLWH worldwide.

## Acknowledgements

We would like to thank all the participants in the study for their time and effort, the POPPY and AGE<sub>h</sub>IV study teams at their respective sites and all the people involved in the COBRA collaboration as detailed below.

The COBRA Steering Committee: P. Reiss (chair), A. Winston, F.W. Wit, M. Prins, M.F. Schim van der Loeff, J. Schouten, B. Schmand, G.J. Geurtsen, D.J. Sharp, M.W.A. Caan, C. Majoie, J. Villaudy, B. Berkhout, N.A. Kootstra, M. Gisslén, A. Pasternak, C.A. Sabin, G. Guaraldi, A. Bürkle, C. Libert, C. Franceschi, A. Kalsbeek, E. Fliers, J. Hoeijmakers, J. Pothof, M. van der Valk, P.H. Bisschop, P. Portegies, S. Zaheri and D. Burger.

The COBRA Project Management Board: P. Reiss, A. Winston, F.W. Wit, J.H. Cole, M.W.A. Caan, J. Villaudy, N.A. Kootstra, M.F. Schim van der Loeff, M. Gisslén, C.A. Sabin, A. Bürkle and W. Zikkenheiner.

The Management Team: P. Reiss, W. Zikkenheiner, F.W. Wit, F.R. Janssen.

The Clinical Cohort Team: A. Winston, F.W. Wit, J. Underwood, J. Schouten, K.W. Kooij, R.A. van Zoest, N. Doyle, M. Prins, M. Schim van der Loeff, P. Portegies, B.A. Schmand, G.J. Geurtsen, E. Verheij, S.O. Verboeket, B.C. Elsenga, M. van der Valk, S. Zaheri, M.M.J. Hillebregt, Y.M.C. Ruijs, D.P. Benschop, L. Tembo, L. McDonald, M. Stott, K. Legg, A. Lovell, O. Erlwein, C. Kingsley, P. Norsworthy, S. Mullaney, T. Kruijer, L. del Grande, V. Olthof, G.R. Visser, L. May, F. Verbraak, N. Demirkaya, I. Visser, G. Guaraldi.

The Neuroimaging Team: D.J. Sharp, M.W.A. Caan, J.H. Cole, C.B.L.M. Majoie, T. Su, R. Leech, J. Huguet.

The HIS Mouse Study Team: J. Villaudy, E. Frankin, A. Pasternak, B. Berkhout, A. van der Kuyl, K. Weijer, E. Siteur-Van Rijnstra.

The Biomarker Team: N.A. Kootstra, M. Gisslén, A.M. Harskamp-Holwerda, I. Maurer, M.M. Mangas Ruiz, A.F. Girigorie, B. Boeser-Nunnink, A. Kalsbeek, P.H.L.T. Bisschop, D. Burger, M. de Graaff-Teulen, J. Hoeijmakers, J. Pothof, C. Libert, S. Dewaele, C. Franceschi, P. Garagnani, C. Pirazzini, M. Capri, F. Dall'Olio, M. Chiricolo, S. Salvioli, D. Fuchs, H. Zetterberg, D. Weber, T. Grune, E.H.J.M. Jansen.

The Data Management and Analysis Team: C.A. Sabin, D. De Francesco, F.W. Wit.

The Dissemination Team: A. Bürkle, T. Sindlinger, S. Oehlke, W. Zikkenheiner, R.A. van Zoest.

This project has received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement n° 305522.

Authors' contributions: D.D.F. contributed to study concept and design, literature search, data analysis and interpretation, figures, writing of the manuscript. F.W.W., N.A.K., A.W., C.A.S., P.R. contributed to study concept and design, data collection and interpretation, critical revision of manuscript. A.B. and S.O. contributed to data analysis and interpretation, critical revision of manuscript. C.F., P.G., C.P., C.L., T.G., D.W., E.H.J.M.J. contributed to data collection, critical revision of manuscript. All authors read and approved the final manuscript.



Source of funding: This study was supported by a European Union Seventh Framework Programme grant to the Comorbidity in Relation to AIDS (COBRA) project (FP-7-HEALTH 305522, all authors), National Institute for Health Research (NIHR) Professorship (NIHR-RP-011-048; DJS), NIHR Imperial Biomedical Research Centre, the Netherlands Organisation for Health Research and Development (ZonMW) (grant number 300020007) & Stichting AIDS Fonds (grant number 2009063), Nuts-Ohra Foundation (grant number 1003-026) and unrestricted scientific grants from: Gilead Sciences, ViiV Healthcare, Janssen Pharmaceutica N.V. Bristol-Myers Squibb (BMS), and Merck & Co to the AGEHIV cohort study (PI Professor P.R.), as well as investigator initiated grants from BMS, Gilead Sciences, Janssen, Merck and ViiV Healthcare to the POPPY cohort study (PI's Professor A.W and Professor C.S.). Funders had no role in analyzing the data, writing the manuscript or in deciding to submit the manuscript for publication.

## Conflicts of interest

There are no conflicts of interest.

## References

- Schouten J, Wit FW, Stolte IG, Kootstra N, van der Valk M, Geerlings SG, et al., AGEHIV Cohort Study Group. **Cross-sectional comparison of the prevalence of age-associated comorbidities and their risk factors between HIV-infected and uninfected individuals: the AGEHIV cohort study.** *Clin Infect Dis* 2014; **59**:1787–1797.
- Althoff KN, McGinnis KA, Wyatt CM, Freiberg MS, Gilbert C, Oursler KK, et al. **Comparison of risk and age at diagnosis of myocardial infarction, end-stage renal disease, and non-AIDS-defining cancer in HIV-infected versus uninfected adults.** *Clin Infect Dis* 2014; **60**:627–638.
- Guaraldi G, Orlando G, Zona S, Menozzi M, Carli F, Garlassi E, et al. **Premature age-related comorbidities among HIV-infected persons compared with the general population.** *Clin Infect Dis* 2011; **53**:1120–1126.
- High KP, Brennan-Ing M, Clifford DB, Cohen MH, Currier J, Deeks SG, et al. **HIV and aging: state of knowledge and areas of critical need for research. A report to the NIH Office of AIDS Research by the HIV and Aging Working Group.** *J Acquir Immune Defic Syndromes* 2012; **60** (Suppl 1): S1–18.
- Simm A, Nass N, Bartling B, Hofmann B, Silber R-E, Navarrete Santos A. **Potential biomarkers of ageing.** *Biol Chem* 2008; **389**:257–265.
- Kennedy BK, Berger SL, Brunet A, Campisi J, Cuervo AM, Epel ES, et al. **Geroscience: linking aging to chronic disease.** *Cell* 2014; **159**:709–713.
- Deelen J, Beekman M, Capri M, Franceschi C, Slagboom PE. **Identifying the genomic determinants of aging and longevity in human population studies: progress and challenges.** *Bioessays* 2013; **35**:386–396.
- Baker GT, Sprott RL. **Biomarkers of aging.** *Expl Gerontol* 1988; **23**:223–239.
- Lagathu C, Cossarizza A, Bérézziat V, Nasi M, Capeau J, Pinti M. **Basic science and pathogenesis of ageing with HIV: potential mechanisms and biomarkers.** *AIDS* 2017; **31**:S105–S119.
- Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, et al. **Microbial translocation is a cause of systemic immune activation in chronic HIV infection.** *Nat Med* 2006; **12**:1365–1371.
- Burdo TH, Lentz MR, Autissier P, Krishnan A, Halpern E, Letendre S, et al. **Soluble CD163 made by monocyte/macrophages is a novel marker of HIV activity in early and chronic infection prior to and after antiretroviral therapy.** *J Infect Dis* 2011; **204**:154–163.
- Dolan SE, Hadigan C, Killilea KM, Sullivan MP, Hemphill L, Lees RS, et al. **Increased cardiovascular disease risk indices in HIV-infected women.** *JAIDS J Acquir Immune Defic Syndr* 2005; **39**:44–54.
- Duprez DA, Neuhaus J, Kuller LH, Tracy R, Belloso W, De Wit S, et al. **Inflammation, coagulation and cardiovascular disease in HIV-infected individuals.** *PLoS One* 2012; **7**:e44454.
- Pathai S, Lawn SD, Gilbert CE, McGuinness D, McGlynn L, Weiss HA, et al. **Accelerated biological ageing in HIV-infected individuals in South Africa: a case-control study.** *AIDS* 2013; **27**:2375–2384.
- Payne BA, Wilson IJ, Hateley CA, Horvath R, Santibanez-Koref M, Samuels DC, et al. **Mitochondrial aging is accelerated by antiretroviral therapy through the clonal expansion of mtDNA mutations.** *Nat Genet* 2011; **43**:806–810.
- Horvath S, Levine AJ. **HIV-1 infection accelerates age according to the epigenetic clock.** *J Infect Dis* 2015; **212**:1563–1573.
- Pathai S, Shiels PG, Weiss HA, Gilbert CE, Peto T, Bekker L-G, et al. **Ocular parameters of biological ageing in HIV-infected individuals in South Africa: relationship with chronological age and systemic biomarkers of ageing.** *Mechanisms Ageing Dev* 2013; **134**:400–406.
- Cole JH, Underwood J, Caan MWA, De Francesco D, van Zoest RA, Leech R, et al., COBRA collaboration. **Increased brain-predicted ageing in treated HIV disease.** *Neurology* 2017; **88**:1349–1357.
- Garagnani P, Bacalini MG, Pirazzini C, Gori D, Giuliani C, Mari D, et al. **Methylation of ELOVL2 gene as a new epigenetic marker of age.** *Aging Cell* 2012; **11**:1132–1134.
- Vanhooren V, Laroy W, Libert C, Chen C. **N-Glycan profiling in the study of human aging.** *Biogerontology* 2008; **9**:351–356.
- Bürkle A, Moreno-Villanueva M, Bernhard J, Blasco M, Zondag G, Hoesjmakers JH, et al. **MARK-AGE biomarkers of ageing.** *Mech Ageing Dev* 2015; **151**:2–12.
- Bürkle A, Berthold M, Junk M, Moreno-Villanueva M. Method for the determination of biological age in human beings. Google Patents; 2016.
- De Francesco D, Wit FW, Cole JH, Kootstra NA, Winston A, Sabin CA, et al., COmorBidity in Relation to AIDS (COBRA) collaboration. **The 'COmorBidity in Relation to AIDS' (COBRA) cohort: design, methods and participant characteristics.** *PLoS One* 2018; **13**:e0191791.
- Weber D, Stuetz W, Bernhard W, Franz A, Raith M, Grune T, et al. **Oxidative stress markers and micronutrients in maternal and cord blood in relation to neonatal outcome.** *Eur J Clin Nutr* 2014; **68**:215–222.
- Jansen E, Beekhof P, Cremers J, Weinberger B, Fiegl S, Toussaint O, et al. **Quality control data of physiological and immunological biomarkers measured in serum and plasma.** *Mech Ageing Dev* 2015; **151**:54–59.
- Bacalini MG, Deelen J, Pirazzini C, De Cecco M, Giuliani C, Lanzarini C, et al. **Systemic age-associated DNA hypermethylation of ELOVL2 gene: in vivo and in vitro evidences of a cell replication process.** *J Gerontol A Biol Sci Med Sci* 2017; **72**:1015–1023.
- Korndewal MJ, Mollema L, Tcherniaeva I, van der Klis F, Kroes ACM, Oudesluys-Murphy AM, et al. **Cytomegalovirus infection in the Netherlands: seroprevalence, risk factors, and implications.** *J Clin Virol* 2015; **63**:53–58.
- Brunner S, Herndler-Brandstetter D, Weinberger B, Grubeck-Loebenstien B. **Persistent viral infections and immune aging.** *Ageing Res Rev* 2011; **10**:362–369.
- Parrinello CM, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ, et al. **Cytomegalovirus immunoglobulin G antibody is associated with subclinical carotid artery disease among HIV-infected women.** *J Infect Dis* 2012; **205**:1788–1796.

30. Gkrania-Klotsas E, Langenberg C, Sharp SJ, Luben R, Khaw K-T, Wareham NJ. **Higher immunoglobulin G antibody levels against cytomegalovirus are associated with incident ischemic heart disease in the population-based EPIC-Norfolk cohort.** *J Infect Dis* 2012; **206**:1897–1903.
31. Booiman T, Wit FW, Girigorie AF, Maurer I, De Francesco D, Sabin CA, et al. **Terminal differentiation of T cells is strongly associated with CMV infection and increased in HIV-positive individuals on ART and lifestyle matched controls.** *PLoS one* 2017; **12**:e0183357.
32. Naggie S. **Hepatitis C virus, inflammation, and cellular aging: turning back time.** *Top Antivir Med* 2017; **25**:3–26.
33. Trickey A, May MT, Schommers P, Tate J, Ingle SM, Guest JL, et al. **CD4: CD8 ratio and CD8 count as prognostic markers for mortality in human immunodeficiency virus-infected patients on antiretroviral therapy: the Antiretroviral Therapy Cohort Collaboration (ART-CC).** *Clin Infect Dis* 2017; **65**:959–966.
34. Bauer J, Baliga R, Liu C, Jones C, Hoyt D. **Endothelial toxicity induced by HIV protease inhibitors: evidence of oxidant related dysfunction and apoptosis.** *Pediatr Res* 2002; **51**:1465A–1465A.
35. Afonso P, Auclair M, Boccarda F, Vantighem M-C, Katlama C, Capeau J, et al. **LMNA mutations resulting in lipodystrophy and HIV protease inhibitors trigger vascular smooth muscle cell senescence and calcification: Role of ZMPSTE24 downregulation.** *Atherosclerosis* 2016; **245**:200–211.
36. Jiménez VC, Wit FW, Joerink M, Maurer I, Harskamp AM, Schouten J, et al. **T-cell activation independently associates with immune senescence in HIV-infected recipients of long-term antiretroviral treatment.** *J Infect Dis* 2016; **214**:216–225.
37. Chou JP, Ramirez CM, Wu JE, Effros RB. **Accelerated aging in HIV/AIDS: novel biomarkers of senescent human CD8+ T cells.** *PLoS One* 2013; **8**:e64702.
38. Gross AM, Jaeger PA, Kreisberg JF, Licon K, Jepsen KL, Khosroheidari M, et al. **Methylome-wide analysis of chronic HIV infection reveals five-year increase in biological age and epigenetic targeting of HLA.** *Mol cell* 2016; **62**:157–168.
39. Pathai S, Lawn SD, Shiels PG, Weiss HA, Cook C, Wood R, et al. **Corneal endothelial cells provide evidence of accelerated cellular senescence associated with HIV infection: a case-control study.** *PLoS One* 2013; **8**:e57422.
40. Pathai S, Lawn SD, Weiss HA, Cook C, Bekker L-G, Gilbert CE. **Increased ocular lens density in HIV-infected individuals with low nadir CD4 counts in South Africa: evidence of accelerated aging.** *J Acquir Immune Defic Syndr* 2013; **63**:307–314.
41. Cevenini E, Invidia L, Lescai F, Salvioli S, Tieri P, Castellani G, et al. **Human models of aging and longevity.** *Expert Opin Biol Ther* 2008; **8**:1393–1405.
42. Sabin CA, Reiss P. **Epidemiology of ageing with HIV: what can we learn from cohorts?** *AIDS* 2017; **31** (Suppl 2):S121–S128.
43. Moreno-Villanueva M, Bürkle A. **Molecular consequences of psychological stress in human aging.** *Exp Gerontol* 2015; **68**:39–42.