

Diagnosing and Managing Diabetes in HIV-Infected Patients: Current Concepts

Anne K. Monroe,¹ Marshall J. Glesby,² and Todd T. Brown³

¹Division of General Internal Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland; ²Division of Infectious Diseases, Weill Cornell Medical College, New York, New York; and ³Division of Endocrinology, Diabetes, and Metabolism, Johns Hopkins University School of Medicine, Baltimore, Maryland

Diabetes mellitus (DM) is a common condition with significant associated morbidity and mortality. DM diagnosis and management among human immunodeficiency virus (HIV)-infected patients is a particularly relevant topic as the HIV-infected population ages and more HIV-infected individuals live with chronic medical comorbidities. Although there is mixed evidence regarding HIV as an independent risk factor for DM, multiple factors related to HIV and its treatment are associated with DM. This review covers the epidemiology of DM in HIV-infected patients, and diagnosis, management, and treatment goals for DM in HIV-infected patients. We highlight the most recent DM treatment guidelines from the American Diabetes Association and the European Association for the Study of Diabetes, emphasizing individualization of DM medication therapy and treatment goals. Finally, we review a comprehensive approach to cardiovascular disease risk reduction in HIV-infected patients with DM and measures to prevent other complications of DM.

Keywords. HIV; diabetes mellitus; CVD.

Diabetes mellitus (DM) is a common condition, affecting 8.5% of the US population [1], with DM prevalence estimates in human immunodeficiency virus (HIV)-infected patients of up to 14% [2]. DM is a leading cause of cardiovascular disease (CVD), blindness, end-stage renal disease, amputations, and hospitalizations.

The goals of this review are to update clinicians on the diagnosis and management of DM in HIV-infected patients, with detailed discussion of treatment goals and CVD risk reduction. Although the management of DM in HIV-infected patients follows the same general guidelines as in HIV-uninfected patients, there are specific considerations with respect to the use of hemoglobin A1c (HbA1c) testing in HIV-infected individuals and

the interactions of particular medications with antiretroviral therapy (ART).

EPIDEMIOLOGY OF DM IN HIV-INFECTED PATIENTS

The prevalence of DM in HIV-infected patients has been reported to range from 2% to 14% and varies by the composition of the cohort studied, how DM diagnosis is ascertained, and how DM risk factors are accounted for in the analysis [2–5]. There is conflicting evidence on whether HIV infection is an independent risk factor for DM, with some studies showing increased risk [2, 6, 7] and others showing no independent effect of HIV on DM or showing an inverse effect [4, 8, 9].

Despite the conflicting evidence on the independent role of HIV in DM, certain factors are clearly associated with DM, including increasing age, obesity, and genetic factors. Other factors influence DM incidence in the general population but are more common in HIV-infected patients: hepatitis C virus infection [10], use of certain medications (atypical antipsychotics, corticosteroids), opiate use, and low testosterone [11]. Furthermore, ART-associated lipodystrophy [4] and visceral fat

Received 26 March 2014; accepted 22 September 2014; electronically published 13 October 2014.

Correspondence: Anne K. Monroe, MD, MSPH, Division of General Internal Medicine, Johns Hopkins University School of Medicine, 1830 E Monument St, Rm 8060, Baltimore, MD 21287 (amonroe4@jhmi.edu).

Clinical Infectious Diseases® 2015;60(3):453–62

© The Author 2014. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

DOI: 10.1093/cid/ciu779

accumulation/lipohypertrophy [3, 12] and HIV-related inflammation (increased proinflammatory cytokines and/or free fatty acids) [13, 14] are DM risk factors in HIV-infected patients. For additional information regarding etiology of DM in HIV-infected patients, readers are referred to a comprehensive article [15].

EFFECTS OF SPECIFIC MEDICATIONS ON DM RISK

Antiretroviral Therapy

Preclinical studies demonstrate that protease inhibitors (PIs) increase insulin resistance through effects on the GLUT-4 transporter and decrease insulin secretion through effects on β -cell function, but the clinical impact of these changes is unclear [16]. It is clear, however, that exposure to certain first-generation ART agents, such as stavudine and indinavir, is strongly associated with developing DM. Use of these agents peaked in 1997–1998 and decreased through 2007–2009. Reflecting this, one study revealed DM incidence peaking in 1999–2000 at 23.2 per 1000 person-years of follow-up (PYFU) and decreasing thereafter to the current level of 5 per 1000 PYFU [17]. A head-to-head comparison of currently prescribed ART regimens revealed that efavirenz increased glucose to a greater degree than atazanavir, although the difference was small (about 4 mg/dL) [18], and the overall effect of current ART on glucose metabolism is modest in most patients.

Statins

Statins are commonly used in HIV-infected patients and can increase insulin resistance and DM [19], although results in studies of HIV-infected patients have been mixed [20–22]. Given the cardiovascular (CV) event reduction benefit from statins, increases in insulin resistance/DM likely do not outweigh the benefit of statin therapy in the general population or in HIV-infected patients [23, 24].

DIABETES DIAGNOSIS

Diabetes and Prediabetes Definitions

Table 1 shows the current American Diabetes Association (ADA) definitions of DM and prediabetes [25].

Table 1. Definitions of Prediabetes and Diabetes^a

| | Hemoglobin A1c ^b | Fasting Plasma Glucose | Random Plasma Glucose | Oral Glucose Tolerance Test |
|-------------|-----------------------------|---------------------------------------|---|--|
| Diabetes | $\geq 6.5\%$ | ≥ 126 mg/dL (≥ 7.0 mmol/L) | ≥ 200 mg/dL (≥ 11.1 mmol/L) with polyuria and polydipsia | ≥ 200 mg/dL (≥ 11.1 mmol/L) |
| Prediabetes | 5.7%–6.4% | 100–125 mg/dL (5.6–6.9 mmol/L) | ... | 140–199 mg/dL (7.8–11.1 mmol/L) |
| Normal | $< 5.7\%$ | ≤ 99 mg/dL (≤ 5.5 mmol/L) | ... | ≤ 139 mg/dL (≤ 7.7 mmol/L) |

^a All laboratory values should be confirmed with repeat testing.

^b Hemoglobin A1c may be inaccurate in the setting of human immunodeficiency virus, so it is not recommended by these authors for diabetes diagnosis.

Use of HbA1c in HIV-Infected Patients

HbA1c is the percentage of glycosylated hemoglobin and reflects long-term glucose status. In conditions associated with shortened red blood cell lifespan, such as hemolytic anemia and pregnancy, only fasting plasma glucose (FPG) or oral glucose tolerance testing should be used to diagnose DM [25, 26]. Data are accumulating that HbA1c may underestimate glycemia in HIV-infected individuals. Although the degree of discordance has varied, higher mean corpuscular volume, nucleoside reverse transcriptase inhibitor use (specifically abacavir), and lower CD4 count have been associated with discordance [27–31].

Testing Recommendations

In view of the possible discordance between HbA1c and glycemic control, we believe it would be prudent to use FPG for the diagnosis of DM, particularly in patients who have risk factors for HbA1c inaccuracy [32]. We agree with the Department of Health and Human Services guidelines that FPG testing should be performed every 6–12 months in all HIV-infected patients and clinicians should consider testing 1–3 months after starting ART [33].

PREVENTING DIABETES

Although a detailed review of DM prevention is beyond the scope of this article, readers are referred to the Standards of Medical Care in Diabetes for DM prevention guidelines (page S20; available at: http://care.diabetesjournals.org/content/37/Supplement_1/S14.extract). Practical implementation of these guidelines may be achieved through community-based options, such as the evidence-based Young Men's Christian Association DM prevention program (Available at: <http://www.ymca.net/diabetes-prevention/>).

DIABETES MANAGEMENT

New Guidelines Focused on Individual Management

The current guidelines for DM management released by the ADA and the European Association for the Study of Diabetes in 2012 emphasize a patient-centered approach in which recommendations are “based on the needs, preferences, and tolerances

of each patient” [34]. Because there is a dearth of head-to-head trials of second-line DM medications, management decisions depend on both the preferences of the patient and the desired clinical outcomes.

Initial Management

Lifestyle modification can have a meaningful impact on glucose control and the course of DM. Referral to a registered dietician for medical nutrition therapy is recommended for all patients with DM [25], as even modest weight loss (as little as 2 kg) can have an impact on glycemic control [35, 36]. Additionally, moderate-intensity aerobic physical activity is recommended for adults with DM [25].

Switching ART Regimens

Switching ART regimens should be considered if a patient is on lopinavir/ritonavir or a thymidine analogue (zidovudine, stavudine) [37]; however, switching other medications is of uncertain benefit.

Diet Recommendations

The American Heart Association (AHA)/American College of Cardiology (ACC)/The Obesity Society guidelines for management of overweight and obesity in adults recommend a reduced-calorie intake diet as part of a comprehensive lifestyle modification intervention [38]. Calorie guidelines for weight loss are (1) 1200–1500 calories/day for women or 1500–1800 calories/day for men; (2) an energy deficit of 500 or 750 calories per day, based on the individual; or (3) an evidence-based diet that restricts a certain food type (eg, high-carbohydrate foods) to create an energy deficit. Dietary recommendations for patients with DM include monitoring carbohydrate intake, limiting consumption of sugar-sweetened beverages, and following a Mediterranean-style diet [25]. A patient-friendly diet guide is available at (<http://www.diabetes.org/food-and-fitness/food/what-can-i-eat/>)

Exercise Recommendations

Aerobic exercise is recommended for at least 150 minutes a week, spread out over at least 3 days per week, along with strength training twice a week. Clinicians can recommend several strategies to increase physical activity among their patients [39–43]: an exercise partner, use of a pedometer with a target (eg, 10 000 steps/day), or individualized counseling with exercise prescription. Linking patients with community- or workplace-based programs may increase exercise uptake.

Medication Therapy

If glucose control is not optimized with lifestyle modification alone, medication therapy should be initiated. Table 2 provides a summary of DM medications highlighting special considerations in HIV-infected patients.

Metformin

The first-line medication for DM is metformin. Advantages to metformin include an average 1% decline in HbA1c, a long track record of safety and efficacy data, no hypoglycemia when given alone, no weight gain, and low cost (generic). Additionally, there may be an independent CVD risk reduction benefit to metformin [44]. Disadvantages include gastrointestinal side effects, particularly nausea and diarrhea, which can be minimized by prescribing a low dose and titrating up. Another rare side effect is lactic acidosis. The contraindications to metformin reflect the associated increased risk of lactic acidosis: chronic kidney disease with creatinine >1.4 mg/dL in women and >1.5 mg/dL in men, hypoxia, decompensated liver disease, severe congestive heart failure (CHF), alcohol abuse, and past history of lactic acidosis (unless there is a remote history specifically related to stavudine or didanosine without recurrence). Special caution should be used when metformin is coadministered with dolutegravir, as dolutegravir increases metformin concentration [45].

Second-line Treatments

After lifestyle modification and metformin, if a patient is still not at goal, there are multiple treatment options.

Sulfonylureas

Sulfonylureas (eg, glipizide and glyburide) act by stimulating insulin release from pancreatic β -cells. Advantages of sulfonylureas include a 1% HbA1c decrease, a long track record of safety and efficacy data, proven decreases in microvascular events, and relatively low cost (generic). Disadvantages include weight gain (2–4 kg), hypoglycemia due to the mechanism of action, and high failure rate (ie, a short time of control on sulfonylurea before additional therapy is required).

Thiazolidinediones

The thiazolidinediones (rosiglitazone and pioglitazone) work by improving target cell response to insulin, thereby lowering glucose. The advantages of thiazolidinedione treatment are a 1% HbA1c decrease and no hypoglycemia. There is a potential independent CVD risk reduction benefit of pioglitazone [46], and pioglitazone raises high-density lipoprotein, lowers triglycerides, and decreases liver fat [47]. There is a low failure rate of thiazolidinediones compared with sulfonylureas, and a modest beneficial effect on lipotrophy may be achieved [48]. Disadvantages include high cost (no generic formulation available), weight gain, fluid retention, and worsening CHF (including in patients with diastolic dysfunction), macular edema, osteoporosis/fracture, and possibly increased risk of bladder cancer [49, 50]. Sales of rosiglitazone were previously restricted due to concern for increased cardiovascular events, however, after US Food and Drug Administration review, the restrictions were removed in mid-2013.

Table 2. Oral Diabetes Medications With Special Considerations in HIV

| Class | Mechanism | Generic Name | Brand Name | Special Considerations in HIV |
|--------------------|--|---|--|--|
| Biguanides | <ul style="list-style-type: none"> Decrease hepatic glucose production Improve insulin sensitivity | Metformin | Glucophage, Glucophage XR, Fortamet, Glumetza, Riomet | <ul style="list-style-type: none"> Dolutegravir increases metformin concentrations |
| Sulfonylureas | <ul style="list-style-type: none"> Stimulate insulin release from the pancreatic β cells Reduce glucose output from the liver Increase insulin sensitivity | Glimepiride Glyburide Micronized glyburide Glipizide | Amaryl Diabeta, Micronase Glynase Glucotrol, Glucotrol xl | |
| Thiazolidinediones | <ul style="list-style-type: none"> Lower blood glucose by improving target cell response to insulin Do not increase pancreatic insulin secretion | Rosiglitazone Pioglitazone | Avandia Actos | <ul style="list-style-type: none"> When used with CYP2C8 inhibitors (many PIs), rosiglitazone/pioglitazone levels may increase. Monitor carefully. |
| Incretins | | | | |
| GLP-1 Analogues | <ul style="list-style-type: none"> Increase glucose-dependent insulin secretion Decrease inappropriate glucagon secretion Increase β-cell growth/replication Slow gastric emptying | Liraglutide Exenatide Exenatide LAR | Victoza Byetta Byetta LAR | |
| DPP-IV inhibitors | <ul style="list-style-type: none"> Increase insulin synthesis and release from pancreatic β cells Decrease glucagon secretion from pancreatic α cells Decreased glucagon secretion results in decreased hepatic glucose production | Sitagliptin Saxagliptin Vildagliptin Linagliptin | Januvia Onglyza Galvus Tradjenta | <ul style="list-style-type: none"> Gliptins have molecular targets on immune cells; however, no evidence that gliptin use changes CD4 or HIV RNA Saxagliptin interacts with strong CYP3A4/5 inhibitors (eg, ritonavir); reduce saxagliptin dose when used with CYP3A4/5 inhibitors |
| Gliflozins | <ul style="list-style-type: none"> Reduce reabsorption of filtered glucose from the tubular lumen and lower the renal threshold for glucose Increase urinary excretion of glucose | Dapagliflozin Canagliflozin | Farxiga Invokana | <ul style="list-style-type: none"> If UGT enzyme inducers (eg, ritonavir) are coadministered with canagliflozin, consider increasing the dose to 300 mg |
| Meglitinides | <ul style="list-style-type: none"> Stimulate insulin release from pancreatic β-cells | Repaglinide Nateglinide | Prandin Starlix | <ul style="list-style-type: none"> When used with CYP3A4/CYP2C8 inhibitors (many PIs), Repaglinide/nateglinide levels may increase. Monitor carefully. EFV and ETR may increase nateglinide |

Abbreviations: CYP, cytochrome P450; DPP-IV, dipeptidyl peptidase 4; EFV, efavirenz; ETR, etravirine; GLP-1, glucagon-like peptide 1; HIV, human immunodeficiency virus; LAR, long-acting release; PI, protease inhibitor; UGT, UDP-glucuronosyltransferase.

Insulin

Insulin is the preferred second-line medication for patients with HbA1c $\geq 8.5\%$. The major advantage of insulin is that it can produce major (ie, unlimited) reductions in HbA1c. However, insulin can cause hypoglycemia, it is associated with weight gain, and the insulin analogues (eg, glargine or detemir insulin) are costly. Recently, concern was raised for mitogenic effects of long-acting insulin due to binding with the insulin-like growth factor 1 receptor. However, several large studies showed no effect of long-acting insulin on cancer incidence [51].

Starting Insulin

Insulin therapy can be initiated with bedtime insulin glargine, insulin detemir, or insulin Neutral Protamine Hagedorn at a dose of 10–15 units, with a dose increase of 2–3 units every 3 days until the fasting glucose is <120 mg/dL (6.7 mmol/L) [52]. Adding prandial short-acting insulin (now available in both injected and inhaled forms) [53] may be beneficial if the fasting glucose goal is not achieved with long-acting insulin alone. Insulin is recommended as first-line therapy for patients with HbA1c $>9\%$, severe liver disease, or severe kidney disease [34].

Incretins

Glucagon-like peptide 1 (GLP-1) regulates glucose homeostasis by decreasing β -cell workload and improving β -cell response. After eating, GLP-1 is secreted into the circulation by the L cells (neuroendocrine cells) of the distal ileum and colon. In the β cells, GLP-1 enhances glucose-dependent insulin secretion. GLP-1 also regulates gastric emptying, decreasing insulin demand, decreases postprandial glucagon secretion, and increases satiety [54]. There are two available therapeutic classes that influence GLP-1: GLP-1 analogues and dipeptidyl peptidase 4 (DPP-IV) inhibitors. GLP-1 analogues have the same effects as endogenous GLP-1 but a longer half-life, and DPP-IV inhibitors block the DPP-IV enzyme, which breaks down GLP-1.

Currently available GLP-1 analogues include exenatide, liraglutide, and exenatide long-acting release. The advantages to GLP-1 analogues are a 1% decrease in HbA1c, no hypoglycemia, weight loss, and preservation of β -cell mass/function. Their disadvantages include gastrointestinal side effects (primarily nausea) and high cost. Postmarketing trials are under way to examine possible increased pancreatitis, thyroid cancer, and pancreatic cancer risk [55]. Currently available DPP-IV inhibitors include sitagliptin, saxagliptin, vildagliptin, and linagliptin. Their advantages are similar to those of GLP-1 analogues. Disadvantages include a relatively small reduction in HbA1c (0.5%), gastrointestinal side effects (decreased intensity compared with GLP-1 analogues), possible hypersensitivity reaction, and high cost. Although a possible CVD risk reduction benefit with gliptins was reported, the benefit was not shown in subsequent studies, with one study reporting an increased rate of hospitalization for heart failure [56–58]. Concern regarding gliptin use in HIV-infected individuals was raised, as gliptins have molecular targets on immune cells; however, a small study revealed no changes in CD4 or HIV RNA among treated HIV-infected patients taking sitagliptin [59]. Of note, saxagliptin interacts with strong cytochrome P450 3A4/5 inhibitors (eg, ritonavir), and saxagliptin dose should be reduced when used in combination [60].

Considering the different second-line options, many clinicians would choose a sulfonylurea due to the high efficacy at low cost with a long track record of safety and efficacy. A large National Institute of Diabetes and Digestive and Kidney Diseases–funded trial comparing the sulfonylureas, the incretins (both a DPP-IV inhibitor and a GLP-1 agonist), and long-acting insulin is under way [61]. Currently, decisions regarding second-line (and subsequent) therapy should be individualized, incorporating patients' clinical characteristics and preferences.

ADDITIONAL AGENTS

Gliflozins

Glucose is freely filtered in glomeruli, then reabsorbed in the proximal tubule by the sodium glucose co-transporter 2

(SGLT2). SGLT2 inhibitors such as dapagliflozin and canagliflozin block the reabsorption of glucose in an insulin-dependent fashion, and glucose is then excreted in the urine. The advantages of gliflozins include weight loss (approximately 2 kg), lower blood pressure (BP), and no hypoglycemia. The disadvantages include consequences of glycosuria (urinary tract infection and genital fungal infections), and potentially increased cardiovascular risk, particularly within the first 30 days of initiating canagliflozin [62]. Data from postmarketing studies further exploring this risk are still pending. No interactions between ART and dapagliflozin are expected; however, if UDP-glucuronosyltransferase enzyme inducers (eg, ritonavir) must be coadministered with canagliflozin, clinicians could consider increasing the dose to 300 mg [63].

Meglitinides

This class of medications includes repaglinide (Prandin) and nateglinide (Starlix), which are short-acting medications that increase insulin release. They are dosed immediately prior to a meal, and offer the advantages of less hypoglycemia than the sulfonylureas and more flexibility with dosing. However, they are expensive, require frequent dosing, and only have a modest effect on HbA1c. For these reasons, they are not frequently used.

GLYCEMIC TARGETS

The standard glycemic target for patients with DM is HbA1c <7%, representing a mean blood glucose <150 mg/dL (8.3 mmol/L), fasting and preprandial values <130 mg/dL (7.2 mmol/L), and postprandial glucose <180 mg/dL (10 mmol/L) [34]. In monitoring DM therapy, HIV clinicians should consider that the HbA1c goal may need to be more stringent in HIV-infected patients, reflecting the underestimation of glycemia by HbA1c in HIV. Achieving glycemic targets, particularly in the setting of aggressive therapy in newly diagnosed diabetes [64], may decrease macrovascular complications of DM (coronary artery disease, cerebrovascular disease, and peripheral vascular disease), although study results have been mixed [65–67].

Achieving glycemic targets has been shown to decrease microvascular (retinopathy, neuropathy, and nephropathy) complications of DM [64, 66, 67]. However, adverse consequences of more intensive glucose control have been observed in clinical trials, including increased severe hypoglycemia and increased death [68, 69]. This underscores the need for individualization of targets. For example, patients with low HbA1c at baseline (ie, mild DM) tended to do better with more intensive targets for DM control. Furthermore, tighter control (HbA1c 6.0%–6.5%) is more appropriate for younger, healthier patients, whereas looser control (HbA1c 7.5%–8.0%) may be more

Table 3. Drug Interactions Between Statins and Antiretroviral Therapy

| Statin | Interactions |
|--------------|---|
| Atorvastatin | <p>PI</p> <p>Significant ↑ in atorvastatin levels with most PIs. TPV/r: contraindicated. For other PIs, Start with lowest dosage (10 mg). Monitor antilipid activity and titrate the statin dosage cautiously.</p> <p>NNRTI</p> <p>EFV: ↓ in atorvastatin levels; may need ↑ dosage. ETR: ↓ in atorvastatin levels; may need ↑ dosage. RPV: no dosage adjustments.</p> <p>Integrase inhibitor</p> <p>EVG/cobicistat/TDF/FTC: ↑ atorvastatin possible. Start with lowest dosage. Monitor response and titrate dosage cautiously.</p> |
| Fluvastatin | <p>PI</p> <p>Not well studied; no known significant interactions with most PIs.</p> <p>NNRTI</p> <p>ETR: may ↑ fluvastatin. Monitor response and titrate dosage cautiously.</p> |
| Lovastatin | <p>PI</p> <p>Substantial ↑ in statin levels, high risk of adverse effects. Contraindicated.</p> <p>NNRTI</p> <p>ETR, NVP: ↓ lovastatin possible. Monitor response.</p> <p>Integrase inhibitor</p> <p>EVG/cobicistat/TDF/FTC: significant ↑ lovastatin expected; contraindicated.</p> |
| Pitavastatin | <p>PI</p> <p>No dosage adjustments necessary.</p> <p>NNRTI</p> <p>EFV: No dosage adjustments necessary.</p> |
| Pravastatin | <p>PI</p> <p>Variable effects; moderate ↑ pravastatin AUC and C_{max} with most. No dosage adjustment of pravastatin is required. Exceptions: DRV/r: Consider alternative statin. If prescribed, use lowest possible dosage, monitor carefully. SQV + RTV: ↓ pravastatin AUC. May need to ↑ pravastatin dosage to reach lipid goals.</p> <p>NNRTI</p> <p>EFV: ↓ pravastatin AUC May need to ↑ pravastatin dosage to reach lipid goals.</p> |
| Rosuvastatin | <p>PI</p> <p>ATV/r: ↑ rosuvasatin C_{max}. Consider alternative statin. If prescribed, use lowest possible dosage (5 mg/day), monitor carefully. Do not exceed 10 mg/day. DRV/r: ↑ rosuvasatin C_{max}. Use lowest possible dosage, monitor carefully. LPV/r: ↑ rosuvasatin C_{max}. Consider alternative statin. If prescribed, use lowest possible dosage, monitor carefully. Do not exceed 10 mg/day. TPV/r: ↑ rosuvasatin C_{max}. Use lowest possible dosage, monitor carefully.</p> <p>Integrase inhibitor</p> <p>EVG/cobicistat/TDF/FTC: rosuvasatin AUC ↑ 38%, C_{max} ↑ 89%; titrate statin dose accordingly.</p> |
| Simvastatin | <p>PI</p> <p>Substantial ↑ in simvastatin levels, high risk of adverse effects. Contraindicated.</p> <p>NNRTI</p> <p>EFV: ↓ simvastatin AUC >50%. May need to ↑ simvastatin dosage to reach lipid goals. ETR, NVP: ↓ simvastatin possible.</p> <p>Integrase inhibitor</p> <p>EVG/cobicistat/TDF/FTC: Significant ↑ simvastatin expected; contraindicated.</p> |

Adapted from Department of Health and Human Services Guidelines for the use of Antiretroviral Agents in HIV-1-infected Adults and Adolescents (Available at: <http://aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf> Tables 18a, 18b, and 18d) and Primary Care of Veterans With HIV Manual (section on lipid-lowering medications-ARV interactions); see <http://www.hiv.va.gov/provider/manual-primary-care/lipid-lowering-meds.asp>.

Abbreviations: ↓, decrease; ↑, increase; ARV, antiretroviral; ATV/r, ritonavir-boosted atazanavir; AUC, area under the curve; C_{max} , maximum serum concentration; DRV/r, ritonavir-boosted darunavir; EFV, efavirenz; ETR, etravirine; EVG, elvitegravir; FTC, emtricitabine; LPV/r, ritonavir-boosted lopinavir; NNRTI, nonnucleoside reverse transcriptase inhibitor; NVP, nevirapine; PI, protease inhibitor; RPV, rilpivirine; RTV, ritonavir; SQV, saquinavir; TDF, tenofovir; TPV/r, ritonavir-boosted tipranavir.

appropriate for older patients with multiple comorbidities who are prone to hypoglycemia [34].

MACROVASCULAR ENDPOINT PREVENTION—ADDITIONAL CONSIDERATIONS

CVD risk factors (dyslipidemia, hypertension, smoking, a positive family history of premature coronary disease, and the presence of albuminuria) should be assessed annually. Routine noninvasive CVD screening in asymptomatic patients, with the exception of annual pedal pulse assessment, is not recommended. Peripheral arterial disease screening includes annual assessment of pedal pulses and consideration of ankle-brachial index testing. Screening for cerebrovascular disease is not recommended [25].

Comprehensive CVD Risk Reduction

HIV-infected patients have higher risk of CVD compared with HIV-uninfected controls [6, 70], likely caused a combination of factors including metabolic derangement, lifestyle factors such as smoking, and chronic inflammation. Patients with both HIV and DM are at particularly high risk. Comprehensive interventions have been shown to have a significant impact on CVD among patients with DM [71] and should include:

- A: Antiplatelet therapy
- B: Blood pressure management
- C: Cholesterol management
- D: Diabetes/glucose management
- S: Smoking cessation

Antiplatelet Therapy

Patients with increased CVD risk should receive aspirin (ASA), although ASA resistance, or “lower than normal antiplatelet response to standard doses of aspirin,” is common, reported in up to 57% of general population patients taking aspirin [72], sometimes making ASA ineffective. A pilot study of ASA in HIV-infected patients showed that heightened platelet activation and immune activation in treated HIV disease was attenuated by 1 week of ASA therapy. This is a potential additional benefit of ASA in patients with HIV and DM [73].

Blood Pressure

The Joint National Committee (JNC) 8 guidelines recommend that patients with DM aged ≥ 18 years start pharmacologic therapy to lower BP at systolic blood pressure (SBP) >140 mm Hg or diastolic blood pressure (DBP) >90 mm Hg to treat to goals SBP <140 mm Hg and DBP <90 mm Hg [74]. JNC 8 guidelines recommend that black patients with DM include a thiazide diuretic or calcium channel blocker (CCB) in their antihypertensive regimen and that nonblack diabetic patients include a thiazide diuretic, CCB, angiotensin-converting enzyme inhibitor (ACE), or angiotensin receptor blocker (ARB) in their regimen.

Cholesterol Management

The ACC/AHA cholesterol treatment guidelines recommend that all patients with DM aged 40–75 years with low-density lipoprotein cholesterol (LDL-c) 70–189 mg/dL without clinical atherosclerotic cardiovascular disease take a moderate-intensity (if calculated 10-year CVD risk is $<7.5\%$) or high-intensity (if 10-year calculated CVD risk is $\geq 7.5\%$) statin [75]. These guidelines recommended checking therapeutic response 4–12 weeks after starting therapy and every 3–12 months thereafter. The expected LDL-c reduction is at least 50% with a high-intensity statin and 30%–50% with a medium-intensity statin.

Clinicians should engage in individualized discussions with their patients, taking into consideration that less aggressive treatment and/or targets may be appropriate for certain patients. Multiple drug interactions between statin therapy and antiretroviral therapy exist (Table 3), requiring careful selection and monitoring of therapy [37].

MICROVASCULAR ENDPOINT PREVENTION—ADDITIONAL CONSIDERATIONS

To help prevent microvascular complications, patients should have yearly ophthalmologic exams to detect retinopathy. To avoid nephropathy, good BP and lipid control are crucial, along with a spot urine microalbumin every 12 months to screen for nephropathy as well as ACE inhibitor or ARB treatment in patients with DM with hypertension or microalbuminuria. Patients should have foot exams every 6–12 months and instruction in foot care, with a referral to podiatry if there is evidence of neuropathy (Table 4) [25].

CONCLUSIONS

DM is a prevalent chronic condition with many deleterious effects, which may be accentuated among patients with both DM

Table 4. Prevention of Microvascular Complications in Patients With Diabetes Mellitus

| Target Area | Recommendation |
|-------------|--|
| Retinopathy | Yearly ophthalmologic exam |
| Nephropathy | Optimize blood pressure and lipid control Spot urine microalbumin every 6 mo to screen for nephropathy ACE inhibitor/ARB in patients with hypertension or microalbuminuria |
| Neuropathy | Foot exams every 6–12 mo Instruction in foot care Referral to podiatry if evidence of neuropathy |

Abbreviations: ACE, angiotensin-converting enzyme; ARB, angiotensin receptor blocker.

and HIV. We recommend that clinicians perform regular DM screening in HIV-infected patients; however, we do not recommend using HbA1c for screening. In treating DM, lifestyle changes are critical, as a 5%–10% weight loss can have important metabolic effects. If drug treatment is required, metformin is first-line therapy. Decisions regarding second- and third-line drugs should be individualized. The HbA1c goal is <7% in most patients, but should be individualized depending on a patient's other comorbidities. A comprehensive approach to prevent micro- and macrovascular complications among HIV-infected patients with DM will lead to the best outcomes.

Note

Potential conflicts of interest. All authors: No potential conflicts of interest.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

- Centers for Disease Control and Prevention. Crude and age-adjusted percentage of civilian, noninstitutionalized adults with diagnosed diabetes, United States, 1980–2011. Available at: <http://www.cdc.gov/diabetes/statistics/prev/national/figage.htm>. Accessed 21 March 2014.
- Brown TT, Cole SR, Li X, et al. Antiretroviral therapy and the prevalence and incidence of diabetes mellitus in the multicenter AIDS cohort study. *Arch Intern Med* **2005**; 165:1179–84.
- De Wit S, Sabin CA, Weber R, et al. Incidence and risk factors for new-onset diabetes in HIV-infected patients. *Diabetes Care* **2008**; 31:1224–9.
- Rasmussen LD, Mathiesen ER, Kronborg G, Gerstoft J, Obel N. Risk of diabetes mellitus in persons with and without HIV: a Danish nationwide population-based cohort study. *PLoS One* **2012**; 7:e44575.
- Polsky S, Floris-Moore M, Schoenbaum EE, Klein RS, Arnsten JH, Howard AA. Incident hyperglycaemia among older adults with or at-risk for HIV infection. *Antivir Ther* **2011**; 16:181–8.
- Triant VA, Lee H, Hadigan C, Grinspoon SK. Increased acute myocardial infarction rates and cardiovascular risk factors among patients with human immunodeficiency virus disease. *J Clin Endocrinol Metab* **2007**; 92:2506–12.
- Galli L, Salpietro S, Pellicciotta G, et al. Risk of type 2 diabetes among HIV-infected and healthy subjects in Italy. *Eur J Epidemiol* **2012**; 27:657–65.
- Howard AA, Hoover DR, Anastos K, et al. The effects of opiate use and hepatitis C virus infection on risk of diabetes mellitus in the Women's Interagency HIV Study. *J Acquir Immune Defic Syndr* **2010**; 54:152–9.
- Butt AA, McGinnis K, Rodriguez-Barradas MC, et al. HIV infection and the risk of diabetes mellitus. *AIDS* **2009**; 23:1227–34.
- Mehta SH, Moore RD, Thomas DL, Chaisson RE, Sulkowski MS. The effect of HAART and HCV infection on the development of hyperglycemia among HIV-infected persons. *J Acquir Immune Defic Syndr* **2003**; 33:577–84.
- Monroe AK, Dobs AS, Xu X, et al. Sex hormones, insulin resistance, and diabetes mellitus among men with or at risk for HIV infection. *J Acquir Immune Defic Syndr* **2011**; 58:173–80.
- Ledergerber B, Furrer H, Rickenbach M, et al. Factors associated with the incidence of type 2 diabetes mellitus in HIV infected participants in the Swiss HIV Cohort Study. *Clin Infect Dis* **2007**; 45:111–9.
- Brown TT, Tassiopoulos K, Bosch RJ, Shikuma C, McComsey GA. Association between systemic inflammation and incident diabetes in HIV-infected patients after initiation of antiretroviral therapy. *Diabetes Care* **2010**; 33:2244–9.
- Meininger G, Hadigan C, Laposata M, et al. Elevated concentrations of free fatty acids are associated with increased insulin response to standard glucose challenge in human immunodeficiency virus-infected subjects with fat redistribution. *Metabolism* **2002**; 51:260–6.
- Samaras K. Prevalence and pathogenesis of diabetes mellitus in HIV-1 infection treated with combined antiretroviral therapy. *J Acquir Immune Defic Syndr* **2009**; 50:499–505.
- Hruz PW. Molecular mechanisms for insulin resistance in treated HIV-infection. *Best Pract Res Clin Endocrinol Metab* **2011**; 25:459–68.
- Capeau J, Bouteloup V, Katlama C, et al. Ten-year diabetes incidence in 1046 HIV-infected patients started on a combination antiretroviral treatment. *AIDS* **2012**; 26:303–14.
- Erlandson KM, Kitch D, Tierney C, et al. Impact of randomized antiretroviral therapy initiation on glucose metabolism. *AIDS* **2014**; 28:1451–61.
- Sattar N, Preiss D, Murray HM, et al. Statins and risk of incident diabetes: a collaborative meta-analysis of randomised statin trials. *Lancet* **2010**; 375:735–42.
- McComsey GA, Jiang Y, Erlandson KM, et al. Rosuvastatin improves hip bone mineral density but worsens insulin resistance [CROI abstract 134]. In: Special Issue: Abstracts From the 2014 Conference on Retroviruses and Opportunistic Infections. *Top Antivir Med* **2014**; 22(e-1):34.
- Lichtenstein K, Debes R, Wood K, et al. Statin use is associated with incident diabetes mellitus among patients in the HIV Outpatient Study. In: 20th Conference on Retroviruses and Opportunistic Infections, Atlanta, GA, 3–6 March 2013. Abstract 767.
- Spagnuolo V, Galli L, Poli A, et al. Association between statin use and type-2 diabetes mellitus occurrence among HIV-1+ patients receiving ART. In: 20th Conference on Retroviruses and Opportunistic Infections, Atlanta, GA, 3–6 March 2013. Abstract 766.
- Wang KL, Liu CJ, Chao TF, et al. Statins, risk of diabetes, and implications on outcomes in the general population. *J Am Coll Cardiol* **2012**; 60:1231–8.
- Ridker PM, Pradhan A, MacFadyen JG, et al. Cardiovascular benefits and diabetes risks of statin therapy in primary prevention: an analysis from the JUPITER trial. *Lancet* **2012**; 380:565–71.
- American Diabetic Association. Standards of medical care in diabetes—2014. *Diabetes Care* **2014**; 37(suppl. 1):S14–80.
- Bonora E, Tuomilehto J. The pros and cons of diagnosing diabetes with A1C. *Diabetes Care* **2011**; 34(suppl. 2):S184–90.
- Kim PS, Woods C, Georgoff P, et al. Hemoglobin A1c underestimates glycemia in HIV infection. *Diabetes Care* **2009**; 82:1591–8.
- Diop M-E, Bastard J-P, Meunier N, et al. Inappropriately low glycated hemoglobin values and hemolysis in HIV-infected patients. *AIDS Res Hum Retroviruses* **2006**; 22:1242–7.
- Polgreen PM, Putz D, Stapleton JT. Inaccurate glycosylated hemoglobin A1C measurements in human immunodeficiency virus—positive patients with diabetes mellitus. *Clin Infect Dis* **2003**; 37:e53–6.
- Glesby MJ, Hoover DR, Shi Q, et al. Glycated hemoglobin in diabetic women with and without HIV infection: data from the women's interagency HIV study. *Antivir Ther* **2010**; 15:571–7.
- Slama L, Palella F, Abraham A, et al. Inaccuracy of hemoglobin A1c among HIV-infected: effects of CD4 cell count, antiretroviral therapies, and hematologic parameters. *J Antimicrob Chemother* **2014**; doi:10.1093/jac/dku295.
- Aberg JA, Gallant JE, Ghanem KG, Emmanuel P, Zingman BS, Horberg MA. Primary care guidelines for the management of persons infected with HIV: 2013 update by the HIV Medicine Association of the Infectious Diseases Society of America. *Clin Infect Dis* **2014**; 58:e1–34.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services. Table 3.

- Available at: <http://aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf>. Accessed 25 July 2014.
34. Inzucchi S, Bergenstal R, Buse J, et al. Management of hyperglycaemia in type 2 diabetes: a patient-centered approach. Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* **2012**; 55: 1577–96.
 35. Pi-Sunyer X, Blackburn G, Brancati F, et al. Reduction in weight and cardiovascular disease risk factors in individuals with type 2 diabetes: one-year results of the look AHEAD trial. *Diabetes Care* **2007**; 30: 1374–83.
 36. Esposito K, Maiorino MI, Ciotola M, et al. Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: a randomized trial. *Ann Intern Med* **2009**; 151:306–14.
 37. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services. Tables 14 and 15. Available at: <http://aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf>. Accessed 25 July 2014.
 38. Jensen MD, Ryan DH, Hu FB, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults. *Circulation* **2014**; 129(suppl. 1):S102–38.
 39. Granner ML, Sharpe PA, Hutto BE, Wilcox S, Addy CL. Perceived individual, social, and environmental factors for physical activity and walking. *J Phys Act Health* **2007**; 4:278–93.
 40. Bravata DM, Smith-Spangler C, Sundaram V, et al. Using pedometers to increase physical activity and improve health. *JAMA* **2007**; 298: 2296–304.
 41. Grandes G, Sanchez A, Sanchez-Pinilla RO, et al. Effectiveness of physical activity advice and prescription by physicians in routine primary care: a cluster randomized trial. *Arch Intern Med* **2009**; 169:694.
 42. Mozaffarian D, Afshin A, Benowitz NL, et al. Population approaches to improve diet, physical activity, and smoking habits a scientific statement from the American Heart Association. *Circulation* **2012**; 126:1514–63.
 43. Artinian NT, Fletcher GF, Mozaffarian D, et al. Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults a scientific statement from the American Heart Association. *Circulation* **2010**; 122:406–41.
 44. Lamanna C, Monami M, Marchionni N, Mannucci E. Effect of metformin on cardiovascular events and mortality: a meta-analysis of randomized clinical trials. *Diabetes Obes Metab* **2011**; 13:221–8.
 45. Tivicay (dolutegravir) prescribing information. Available at: http://www.viivhealthcare.com/media/58599/us_tivicay.pdf. Accessed 21 March 2014.
 46. Mannucci E, Monami M, Lamanna C, Gensini G, Marchionni N. Pioglitazone and cardiovascular risk. A comprehensive meta-analysis of randomized clinical trials. *Diabetes Obes Metab* **2008**; 10:1221–38.
 47. Deeg MA, Buse JB, Goldberg RB, et al. Pioglitazone and rosiglitazone have different effects on serum lipoprotein particle concentrations and sizes in patients with type 2 diabetes and dyslipidemia. *Diabetes Care* **2007**; 30:2458–64.
 48. Tungsrirat M, El Bejjani D, Rizk N, et al. Rosiglitazone improves lipoproteins in patients receiving thymidine-sparing regimens. *AIDS* **2010**; 24:1291–8.
 49. Lewis JD, Ferrara A, Peng T, et al. Risk of bladder cancer among diabetic patients treated with pioglitazone interim report of a longitudinal cohort study. *Diabetes Care* **2011**; 34:916–22.
 50. Schernthaner G, Currie CJ, Schernthaner G-H. Do we still need pioglitazone for the treatment of type 2 diabetes? A risk-benefit critique in 2013. *Diabetes Care* **2013**; 36(suppl. 2): S155–61.
 51. Home P, Lagarenne P. Combined randomised controlled trial experience of malignancies in studies using insulin glargine. *Diabetologia* **2009**; 52:2499–506.
 52. Afrezza prescribing information. Available at: <http://www.mannkindcorp.com/product-pipeline-diabetes-afrezza-micro-site.htm>. Accessed 13 October 2014.
 53. Yki-Järvinen H, Juurinen L, Alvarsson M, et al. Initiate Insulin by Aggressive Titration and Education (INITIATE): a randomized study to compare initiation of insulin combination therapy in type 2 diabetic patients individually and in groups. *Diabetes Care* **2007**; 30: 1364–9.
 54. Drucker DJ, Nauck MA. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* **2006**; 368:1696–705.
 55. US Food and Drug Administration. Byetta safety update for healthcare professionals. Available at: <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/DrugSafetyInformationforHealthcareProfessionals/ucm190406.htm>. Accessed 21 March 2014.
 56. Patil HR, Al Badarin FJ, Al Shami HA, et al. Meta-analysis of effect of dipeptidyl peptidase-4 inhibitors on cardiovascular risk in type 2 diabetes mellitus. *Am J Cardiol* **2012**; 110:826–33.
 57. Scirica BM, Bhatt DL, Braunwald E, et al. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* **2013**; 369:1317–26.
 58. White WB, Cannon CP, Heller SR, et al. Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med* **2013**; 369: 1327–35.
 59. Goodwin SR, Reeds DN, Royal M, Struthers H, Laciny E, Yarasheski KE. Dipeptidyl peptidase IV inhibition does not adversely affect immune or virological status in HIV infected men and women: a pilot safety study. *J Clin Endocrinol Metab* **2012**; 98:743–51.
 60. Onglyza (saxagliptin) prescribing information. Available at: http://packageinserts.bms.com/pi/pi_onglyza.pdf. Accessed 21 March 2014.
 61. National Institutes of Health. NIH begins recruitment for long-term study of diabetes drug efficacy. Available at: <http://www.nih.gov/news/health/jun2013/niddk-03.htm>. Accessed 21 March 2014.
 62. Neal B, Perkovic V, de Zeeuw D, et al. Rationale, design, and baseline characteristics of the Canagliflozin Cardiovascular Assessment Study (CANVAS)—a randomized placebo-controlled trial. *Am Heart J* **2013**; 166:217–23.
 63. Invokana (canagliflozin) prescribing information. Available at: <http://www.invokanahcp.com/prescribing-information.pdf>. Accessed 21 March 2014.
 64. Holman RR, Paul SK, Bethel MA, et al. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* **2008**; 359:1577–89.
 65. Abraira C, Colwell J, Nuttall F, et al. Cardiovascular events and correlates in the Veterans Affairs Diabetes Feasibility Trial. Veterans Affairs Cooperative Study on Glycemic Control and Complications in Type II Diabetes. *Arch Intern Med* **1997**; 157:181–8.
 66. Action to Control Cardiovascular Risk in Diabetes Study Group; Gerstein HC, Miller ME, et al. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* **2008**; 358:2545–59.
 67. ADVANCE Collaborative Group; Patel A, MacMahon S, et al. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* **2008**; 358:2560–72.
 68. Kelly TN, Bazzano LA, Fonseca VA, Thethi TK, Reynolds K, He J. Systematic review: glucose control and cardiovascular disease in type 2 diabetes. *Ann Intern Med* **2009**; 151:394–403.
 69. Gerstein HC, Miller ME, Byington RP, et al. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* **2008**; 358:2545–59.
 70. Freiberg MS, Chang CC, Kuller LH, et al. HIV infection and the risk of acute myocardial infarction. *JAMA Intern Med* **2013**; 173: 614–22.
 71. Gaede P, Lund-Andersen H, Parving HH, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* **2008**; 358:580–91.

72. Grinstein J, Cannon CP. Aspirin resistance: current status and role of tailored therapy. *Clin Cardiol* **2012**; 35:673–80.
73. O'Brien M, Montenont E, Hu L, et al. Aspirin attenuates platelet activation and immune activation in HIV-1-infected subjects on antiretroviral therapy: a pilot study. *J Acquir Immune Defic Syndr* **2013**; 63:280–8.
74. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* **2014**; 311:507–20.
75. Stone NJ, Robinson J, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* **2014**; 63(25 pt B):2889–934.

Please excuse the presence of this and the following test pages, which have been added to a small number of article PDFs for a limited time as part of our process of continual development and improvement.

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.