P004

Referral from Post-Exposure Prophylaxis to Pre-Exposure Prophylaxis in the United States, 2016-2025: An Analysis of Real World Claims Data

Supplementary Materials

Kenneth H Mayer[1], Travis Lim[2], Yi-Han Hu[3], Wenyi Wang[2], Laura M Mann[4], Jason T Hindman[2], Lillian B Brown[2]

[1]Fenway Health, Boston, MA, USA; [2]Gilead Sciences, Inc., Foster City, CA, USA; [3]Syneos Health, Morrisville, NC, USA; [4]IQVIA, Atlanta, GA, USA

HIV Drug Therapy Glasgow 2026
8-11 November 2026
Glasgow, United Kingdom

Table S1. ICD-10-CM Diagnosis Codes and LOINC Codes Used to Define Risk Events

Administrative Code[a]	Description
ICD-10-CM diagnosis codes	
Z20.6	Contact with and (suspected) exposure to HIV
Z20.2	Contact with and (suspected) exposure to infections with a predominantly sexual mode of transmission
Z20.5	Contact with and (suspected) exposure to viral hepatitis
Z72.51	High-risk heterosexual behaviour
Z72.52	High-risk homosexual behaviour
Z72.53	High-risk bisexual behaviour
Z77.21	Contact with and (suspected) exposure to potentially hazardous body fluids
Laboratory testing (LOINC) codes	
5287, 450932, 433052, 803668, 803635, 803650	STI test for chlamydia or gonorrhoea based on a rectal swab

[a]ICD-10-CM and LOINC codes were used to define risk exposures linked to PEP episodes. Only ICD-10-CM codes were used to define risk events during the follow-up period.
ICD-10-CM, *International Classification of Diseases, Tenth Revision, Clinical Modification*; LOINC, Logical Observation Identifiers Names and Codes; PEP, post-exposure prophylaxis; STI, sexually transmitted infection.