P-969

Differences in Demographics and HIV Risk Exposures Among People Receiving Bictegravir/Emtricitabine/Tenofovir Alafenamide (B/F/TAF) Versus Other Post-Exposure Prophylaxis Regimens in the United States, 2016-2025

Supplementary Materials

Kenneth H Mayer[1], Travis Lim[2], Valentina Shvachko[2], Laura M Mann[3], Wenyi Wang[2], Jason T Hindman[2]

[1]Fenway Health, Boston, MA, USA; [2]Gilead Sciences, Inc., Foster City, CA, USA; [3]IQVIA, Atlanta, GA, USA

IDWeek
October 21–24, 2026
Washington, DC, USA

Table S1. Risk Exposure Codes

Code Category	Code	Note/Detail	Risk Category
Occupational exposure diagnosis codes	W46.1XXA	Contact with contaminated hypodermic needle	Occupational risk exposure
	Z77.21	Exposure to bodily fluids	Other exposure
Risk exposure diagnosis codes	Z20.2, Z20.5, Z20.6, Z72.51, Z72.52, Z72.53	High-risk sexual behavior, exposure to HIV with sexual transmission	Sexual risk exposure
Laboratory testing (LOINC) codes	57287, 450932, 433052, 803668, 803635, 803650	STI test for chlamydia or gonorrhea	Sexual risk exposure
Sexual assault codes	T74.21XA, Z04.41	Sexual abuse, sexual assault, or rape; adult, initial encounter or examination	Sexual risk exposure

LOINC, Logical Observation Identifiers Names and Codes; STI, sexually transmitted infection.

Table S2. Diagnosis Codes

ICD-10-CM and ICD-9-CM Codes used for HIV Diagnosis	042, V08, 079.53, 795.71, B97.35, R75, B20, Z21, O98.7

ICD-10-CM, *International Classification Of Diseases, Tenth Revision*, Clinical Modification; ICD-9-CM, *International Classification Of Diseases, Ninth Revision*, Clinical Modification.